


HIV Surveillance by Race/Ethnicity

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention



Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity, 2006–2009—40 States and 5 U.S. Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



HIV Infection in Blacks/African Americans— 40 States and 5 U.S. Dependent Areas

Of the 171,840 diagnoses of HIV infection from 2006–2009,
blacks/African Americans accounted for:

50% of total

64% of women

66% of infections attributed to heterosexual^a contact

68% of children aged <13 years

In 2009, 50% of diagnoses of HIV infection among adults and
adolescents were in blacks/African Americans.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



HIV Infection in Hispanics/Latinos — 40 States and 5 U.S. Dependent Areas

Of the 171,840 diagnoses of HIV infection from 2006–2009, Hispanics/Latinos^a accounted for:

20% of total

17% of women

18% of infections attributed to heterosexual contact^b

17% of children aged <13 years

In 2009, 19% of diagnoses of HIV infection among adults and adolescents were in Hispanics/Latinos.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting

^a Hispanics/Latinos can be of any race.


^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



Diagnoses of HIV Infection and Population, by Race/Ethnicity, 2009—40 States


Diagnoses of HIV infection

N=42,011



Population, 40 States

N=241,832,054



 American Indian/Alaska Native

 Asian

 Black/African American

 Multiple races

 Hispanic/Latino^a

 Native Hawaiian/Other Pacific Islander

 White


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.




Diagnoses of HIV Infection and Population among Adult and Adolescent Males, by Race/Ethnicity, 2009—40 States

Diagnoses of HIV Infection
N=31,872



Male Population, 40 States
N=97,416,423



- American Indian/Alaska Native
- Asian
- Black/African American
- Multiple races
- Hispanic/Latino^a
- Native Hawaiian/Other Pacific Islander
- White

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection and Population among Adult and Adolescent Females, by Race/Ethnicity, 2009—40 States


Diagnoses of HIV Infection

N=9,973



Female Population, 40 States

N=102,041,789



 American Indian/Alaska Native

 Asian

 Black/African American

 Multiple races

 Hispanic/Latino^a

 Native Hawaiian/Other Pacific Islander

 White


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.




Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, by Sex and Transmission Category, 2009—40 States and 5 U.S. Dependent Areas

Males
N=14,914



Females
N=6,632



- Male-to-male sexual contact
- Injection drug use
- Male-to-male sexual contact and IDU
- Heterosexual contact^a
- Other^b


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, by Sex and Transmission Category, 2009—40 States and 5 U.S. Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adult and Adolescent Whites, by Sex and Transmission Category, 2009—40 States and 5 U.S. Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.




Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, by Sex and Country of Birth, 2009—40 States and 5 U.S. Dependent Areas

Males
N=6,615



Females
N=1,625



- United States
- Central America
- South America
- Cuba
- Mexico
- Puerto Rico
- Other
- Unknown

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Males, by Race/Ethnicity, 2009—40 States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	137	18.4
Asian	363	12.7
Black/African American	14,898	122.2
Hispanic/Latino ^a	5,972	48.3
Native Hawaiian/Other Pacific Islander	26	41.2
White	10,093	14.8
Multiple races	383	41.3
Total	31,872	32.7

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2009—40 States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	51	6.6
Asian	103	3.4
Black/African American	6,627	47.8
Hispanic/Latino ^a	1,352	11.9
Native Hawaiian/Other Pacific Islander	8	13.3
White	1,699	2.4
Multiple races	132	13.4
Total	9,973	9.8


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, 2009—40 States

N = 21,525




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, 2009—40 States

N = 7,324




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.
^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Whites, 2009—40 States

N = 11,791



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Deaths of Persons with a Diagnosis of HIV Infection, by Race/Ethnicity, 2008—40 States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	71	3.8
Asian ^a	56	0.8
Black/African American	8,845	27.5
Hispanic/Latino ^b	2,565	8.2
Native Hawaiian/Other Pacific Islander	2	1.5
White	4,756	2.9
Multiple races	470	15.7
Total^c	16,769	7.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.



Adults and Adolescents Living with a Diagnosis of HIV Infection, by Race/Ethnicity, Year-end 2008—40 States


Race/ethnicity	No.	Rate
American Indian/Alaska Native	2,373	157.9
Asian ^a	4,196	72.8
Black/African American	313,427	1,216.7
Hispanic/Latino ^b	114,682	498.8
Native Hawaiian/Other Pacific Islander	295	243.4
White	215,366	154.1
Multiple races	8,911	480.3
Total^c	660,061	333.7

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

- ^a Includes Asian/Pacific Islander legacy cases.
- ^b Hispanics/Latinos can be of any race.
- ^c Includes persons of unknown race/ethnicity




AIDS Diagnoses among Minority Races/Ethnicities, 1985-2009, United States and Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Includes American Indians/Alaska Natives, Asians, Blacks/African Americans, Hispanics/Latinos, and Native Hawaiians/Other Pacific Islanders. The Asian category includes Asian/Pacific Islander legacy cases. Hispanics/Latinos can be of any race



AIDS Diagnoses among Adults and Adolescents, by Race/Ethnicity and Year of Diagnosis, 1985-2009— United States and Dependent Areas





Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



AIDS Diagnoses and Population, by Race/Ethnicity, 2009—United States



- American Indian/Alaska Native
- Asian
- Black/African American
- Multiple races
- Hispanic/Latino^a
- Native Hawaiian/Other Pacific Islander
- White


Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Rates of AIDS Diagnoses among Adult and Adolescent Blacks/African Americans, 2009—United States

N = 16,733




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.



Rates of AIDS Diagnoses among Adult and Adolescent Hispanics/Latinos^a, 2009—United States

N = 6,715




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.

^aHispanics/Latinos can be of any race.



Survival after an AIDS Diagnosis during 1998–2005, by Months Survived and Race/Ethnicity— United States and Dependent Areas



Note. Data exclude persons whose month of diagnosis or month of death is unknown.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.



Adults and Adolescents Living with an AIDS Diagnosis, by Race/Ethnicity, Year-end 2008—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	1,729	92.9
Asian ^a	4,855	43.9
Black/African American	208,503	695.9
Hispanic/Latino ^b	94,125	271.7
Native Hawaiian/Other Pacific Islander	439	125.5
White	163,157	95.8
Multiple races	6,201	218.9
Total^c	479,161	190.9

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.

