


# Proportion of HIV/AIDS Cases and Population among Adolescents 13 to 19 Years of Age, by Race/Ethnicity Diagnosed in 2006—33 States




Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.  
 \*Includes 22 persons of unknown race or multiple races.




# Proportion of HIV/AIDS Cases and Population among Young Adults 20 to 24 Years of Age, by Race/Ethnicity Diagnosed in 2006—33 States

HIV/AIDS cases  
N=3,886\*



Population, 33 states  
N = 13,424,369



White, not Hispanic  
Black, not Hispanic  
Hispanic


Asian/Pacific Islander  
American Indian/  
Alaska Native



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.  
\*Includes 22 persons of unknown race or multiple races.



# Proportion of HIV/AIDS Cases among Adults and Adolescents, by Sex and Age Group Diagnosed in 2006—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis.  
Data from 33 states with confidential name-based HIV infection reporting since at least 2003.  
Data based on person's age at diagnosis.  
Data have been adjusted for reporting delays.



# HIV/AIDS Cases among Male Adolescents and Young Adults, by Transmission Category 2003–2006—33 States

Transmission	13–19 years		20–24 years	
	N	%	N	%
Male-to-male sexual contact	2,286	81	7,756	78
Injection drug use (IDU)	146	5	645	7
Male-to-male sexual contact and IDU	100	4	454	5
High-risk heterosexual contact*	277	10	1,013	10
Other/not identified <sup>†</sup>	9	<1	20	<1
<b>Total</b>	<b>2,817</b>	<b>100</b>	<b>9,888</b>	<b>100</b>

Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.

\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.





# HIV/AIDS Cases among Female Adolescents and Young Adults, by Transmission Category 2003–2006—33 States

Transmission Category	13–19 years		20–24 years	
	N	%	N	%
Injection drug use	233	13	597	14
High-risk heterosexual contact*	1,546	86	3,492	85
Other/not Identified†	15	<1	32	<1
Total	1,793	100	4,121	100

Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis.  
 Data from 33 states with confidential name-based HIV infection reporting since at least 2003.  
 Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.


\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.






# HIV/AIDS Cases among Adolescents 13 to 19 Years of Age, 2006—33 States N=1,373



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays. Data based on person's age at diagnosis.



# Adolescents 13 to 19 Years of Age Living with HIV/AIDS, 2006—33 States N=5,678




Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays. Data based on age as of December 31, 2006.



# Reported AIDS Cases among Adolescents 13 to 19 Years of Age, by Sex, 1985–2006—United States and Dependent Areas

N=6,642




Note. Data based on person's age at diagnosis.





# Reported AIDS Cases among Young Adults 20 to 24 Years of Age, by Sex, 1985–2006—United States and Dependent Areas



N=36,293



Note. Data based on person's age at diagnosis.



# Proportion of AIDS Cases and Population among Adolescents 13 to 19 Years of Age, by Race/Ethnicity Reported in 2006—50 States and DC





White, not Hispanic  
 Black, not Hispanic  
 Hispanic

Asian/Pacific Islander  
 American Indian/  
Alaska Native

Note. Data based on person's age at AIDS diagnosis.  
\*Includes 6 persons of unknown race or multiple races.



# Proportion of AIDS Cases and Population among Young Adults 20 to 24 Years of Age, by Race/Ethnicity Reported in 2006—50 States and DC



White, not Hispanic  
Black, not Hispanic  
Hispanic


Asian/Pacific Islander  
American Indian/  
Alaska Native

Note. Data based on person's age at AIDS diagnosis.  
\* Includes 18 persons of unknown race or multiple races.





# Proportion of AIDS Cases among Adults and Adolescents, by Sex and Age Group, Reported in 2006—United States and Dependent Areas



Note. Data based on person's age at diagnosis.



# AIDS Cases among Male Adolescents and Young Adults, by Transmission Category 2002–2006—United States and Dependent Areas

Transmission category	13–19 years		20–24 years	
	N	%	N	%
Male-to-male sexual contact	736	60	3,983	72
Injection drug use (IDU)	86	7	480	9
Male-to-male sexual contact and IDU	38	3	325	6
High-risk heterosexual*	147	12	706	13
Other/not identified <sup>†</sup>	225	18	58	1
<b>Total</b>	<b>1,231</b>	<b>100</b>	<b>5,552</b>	<b>100</b>

Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>†</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# AIDS Cases among Female Adolescents and Young Adults, by Transmission Category 2002–2006—United States and Dependent Areas

Transmission category	13–19 years		20–24 years	
	N	%	N	%
Injection drug	108	11	416	16
High-risk heterosexual contact*	549	59	2,072	81
Other/not identified †	279	30	62	2
Total	936	100	2,550	100

Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.


\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Reported AIDS Cases among Adolescents 13 to 19 Years of Age, 2006—United States and Dependent Areas

N=581




Note. Data based on person's age at diagnosis.





# Adolescents 13 to 19 Years of Age Reported to be Living with AIDS, 2006—United States and Dependent Areas N=3,116\*



Note. Data based on person's age as of December 31, 2006.  
\*Includes 13 persons whose state of residence is unknown or missing.

