


Estimated Numbers and Percentages of AIDS Cases among Adult and Adolescent Men Who Have Sex with Men 1985–2007—United States and Dependent Areas



Note. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.



Estimated AIDS Cases in Males, Cumulative through 2007—50 States and DC

Of the 1,009,219 AIDS cases in adults and adolescents

80% were in males

60% of cases in males were attributed to
male-to-male sexual contact

4% of cases in males were aged 13–24 years

66% of cases in males aged 13–24 were
attributed to male-to male sexual contact


Of AIDS cases diagnosed during 2007, 47% of cases
in adults and adolescents were attributed to male-to-male



Note. Data have been adjusted for reporting delays and missing risk-factor information.
Data exclude cases among men who had sex with other men and injected drugs.



Estimated Numbers of AIDS Cases and Deaths among Men Who Have Sex with Men 1985–2007—United States and Dependent Areas




Note. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.







Estimated Numbers of AIDS Cases among Adult and Adolescent Men Who Have Sex with Men, by Race/Ethnicity 1985–2007—United States and Dependent Areas




Note. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.

*Hispanics/Latinos can be of any race.

†Includes Asian and Pacific Islander legacy cases.



Percentages of Estimated HIV/AIDS Cases among Adult and Adolescent Men Who Have Sex with Men by Race/Ethnicity, 1985–2007—25 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 25 states with confidential name-based HIV infection reporting since at least 1994. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.


*Hispanics/Latinos can be of any race.

†Includes Asian and Pacific Islander legacy cases.





Percentages of Estimated HIV/AIDS Cases among Adult and Adolescent Men Who Have Sex with Men by Race/Ethnicity 2007—34 States




Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 34 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.

†Includes Asian and Pacific Islander legacy cases.

*Hispanics/Latinos can be of any race.



Estimated Numbers of AIDS Cases among Adult and Adolescent Men Who Have Sex with Men, by Region and Race/Ethnicity, 2007—50 States and DC




Note. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.

†Includes Asian and Pacific Islander legacy cases.

*Hispanics/Latinos can be of any race.




Estimated Numbers of HIV/AIDS Cases among Adult and Adolescent Men Who Have Sex with Men, by Age Group 2004–2007—34 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 34 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.



Estimated Numbers of HIV/AIDS Cases among Men Who Have Sex with Men, Aged 13–24 Years, by Race/Ethnicity 2004–2007—34 States




Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 34 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.

†Includes Asian and Pacific Islander legacy cases.

*Hispanics/Latinos can be of any race.



Percentages of HIV/AIDS Cases among Men Who Have Sex with Men, Aged 13–24 Years by Race/Ethnicity, 2007—34 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 34 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and missing risk-factor information. Data exclude cases among men who had sex with other men and injected drugs.

†Includes Asian and Pacific Islander legacy cases.

*Hispanics/Latinos can be of any race.



