



## HIV and Its Treatment

HIV/AIDS: The Basics

Testing for HIV

Seeing an HIV Health Care Provider

When to Start Anti-HIV Medications

Recommended HIV Treatment Regimens

FDA-Approved Anti-HIV Medications

Treatment Adherence

Following an HIV Treatment Regimen

Is My Treatment Regimen Working?

Changing an HIV Treatment Regimen

HIV Coinfections

Preventing Transmission of HIV

Telephone: 1-800-448-0440  
International: 1-301-315-2816  
E-mail: [contactus@aidsinfo.nih.gov](mailto:contactus@aidsinfo.nih.gov)  
Web: <http://aidsinfo.nih.gov>



## HIV and Its Treatment

---

This series of fact sheets on HIV and its treatment is intended for adults and adolescents infected with HIV, their families, and their friends. The fact sheets include information on the basics of HIV/AIDS, recommended anti-HIV medications, and tips on how to successfully follow an anti-HIV regimen.

The fact sheets are based on information in the *Guidelines for the Use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents*. The Guidelines are developed by the U.S. Department of Health and Human Services (HHS) Panel on Antiretroviral Guidelines for Adults and Adolescents, a working group of the Office of AIDS Research Advisory Council (OARAC). The current Guidelines are available on the AIDSinfo website at <http://aidsinfo.nih.gov/guidelines>.

These fact sheets are not intended as a substitute for the expert advice and care of medical professionals. Individuals seeking HIV/AIDS-related medical advice should consult with a health care provider.

### Table of Contents

---

1. HIV/AIDS: The Basics
2. Testing for HIV
3. Seeing an HIV Health Care Provider
4. When to Start Anti-HIV Medications
5. Recommended HIV Treatment Regimens
6. FDA-Approved Anti-HIV Medications
7. Treatment Adherence
8. Following an HIV Treatment Regimen
9. Is My Treatment Regimen Working?
10. Changing an HIV Treatment Regimen
11. HIV Coinfections
12. Preventing Transmission of HIV

## HIV/AIDS: The Basics

### What is HIV/AIDS?

The **human immunodeficiency virus**, or **HIV**, is the virus that causes HIV infection. During HIV infection, the virus attacks and destroys the infection-fighting **CD4 cells** of the body's immune system. Loss of CD4 cells makes it difficult for the immune system to fight infections.

**Acquired immunodeficiency syndrome**, or **AIDS**, is the most advanced stage of HIV infection.

### How is HIV transmitted?

HIV is transmitted (spread) through the blood, semen, genital fluids, or breast milk of a person infected with HIV. Having **unprotected sex** or sharing drug injection equipment (such as needles and syringes) with a person infected with HIV are the most common ways HIV is transmitted.

You can't get HIV by shaking hands, hugging, or closed-mouth kissing with a person who is infected with HIV. And you can't get HIV from contact with objects such as toilet seats, doorknobs, dishes, or drinking glasses used by a person infected with HIV.

Even though it takes many years for symptoms of HIV to develop, a person infected with HIV can spread the virus at any stage of HIV infection. Detecting HIV early after infection and starting treatment with anti-HIV medications before symptoms of HIV develop can help people with HIV live longer, healthier lives. Treatment can also reduce the risk of **transmission of HIV**.

### What is the treatment for HIV?

**Antiretroviral therapy (ART)** is the recommended treatment for HIV infection. ART involves taking a combination (**regimen**) of three or more anti-HIV medications daily. ART prevents HIV from multiplying and destroying infection-fighting CD4 cells. This helps the body fight off life-threatening infections and cancer.

ART can't cure HIV, but anti-HIV medications help people infected with HIV live longer, healthier lives.

### Can treatment prevent HIV from advancing to AIDS?

Yes. Treatment with anti-HIV medications prevents HIV from multiplying and destroying the immune system. This helps the body fight off life-threatening infections and cancers and prevents HIV from advancing to AIDS.

### Terms Used in This Fact Sheet:

**AIDS:** Acquired immunodeficiency syndrome. AIDS is the most advanced stage of HIV infection. AIDS is diagnosed when a person infected with HIV has a CD4 count of less than 200 cells/mm<sup>3</sup> or has an AIDS-defining condition.

**AIDS-defining condition:** Any one of several illnesses that can lead to a diagnosis of AIDS in a person infected with HIV. AIDS is the most advanced stage of HIV infection.

**Antiretroviral therapy (ART):** The recommended treatment for HIV. ART involves taking a combination of three or more anti-HIV medications from at least two different drug classes every day to control the virus.

**CD4 cells:** Also called T cells or CD4+ T cells. Infection-fighting white blood cells of the immune system. HIV destroys CD4 cells, making it harder for the body to fight infections.

**CD4 count:** The number of CD4 cells in a sample of blood. A CD4 count measures how well the immune system is working.

**HIV:** Human immunodeficiency virus. HIV is a virus that attacks the immune system, putting people infected with HIV at risk for life-threatening infections and cancer. AIDS is the most advanced stage of HIV infection.

**Opportunistic infection:** An infection that occurs more frequently or is more severe in people with weakened immune systems (such as people with HIV or people receiving chemotherapy) than in people with healthy immune systems.

**Regimen:** A combination of three or more anti-HIV medications from at least two different drug classes.

**Transmission of HIV:** The spread of HIV from a person infected with HIV to another person through the infected person's blood, semen, genital fluids, or breast milk.

**Unprotected sex:** Sex without using a condom.

It takes many years, but without treatment, HIV infection can advance to AIDS. A diagnosis of AIDS requires that a person infected with HIV have either:

- A **CD4 count** of less than 200 cells/mm<sup>3</sup>. (The CD4 count of a healthy person ranges from 500 to 1,200 cells/mm<sup>3</sup>.)

### OR

- An **AIDS-defining condition**. (AIDS-defining conditions include **opportunistic infections** and cancers that are life-threatening in a person with HIV. Having an AIDS-

defining condition signals that a person's HIV infection has advanced to AIDS.)

### **What illnesses are considered AIDS-defining conditions?**

The Centers for Disease Control and Prevention (CDC) considers several illnesses AIDS-defining conditions.

*Pneumocystis jiroveci* pneumonia, tuberculosis, and toxoplasmosis are examples of AIDS-defining conditions.

### **For more information:**

Contact an AIDSinfo health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.

## Testing for HIV

### I may have been exposed to HIV. What should I do?

Get tested. The only way to know if you're infected with the virus is to get an HIV test.

Soon after infection with HIV, a person may have flu-like symptoms. But HIV infection isn't diagnosed on the basis of symptoms. Getting tested is the only way to know if you're infected with HIV.

### What is the most common HIV test?

The most common HIV test is the **HIV antibody test**. HIV antibodies are a type of protein the body produces in response to HIV infection. The HIV antibody test checks for HIV antibodies in a person's blood, urine, or fluids from the mouth.

Generally it takes the body about 3 months from the time of infection to produce enough antibodies to be detected by an HIV antibody test. (For some people, it can take up to 6 months.) The time period between infection and the appearance of detectable HIV antibodies is called the **window period**. Because HIV antibodies are not detectable yet, the HIV antibody test isn't useful during the window period.

### What HIV test is used during the window period?

The **plasma HIV RNA test** (also called a **viral load** test) can detect HIV in a person's blood within 9 days of infection, **before** the body develops detectable HIV antibodies. The plasma HIV RNA test is recommended when recent infection is very likely—for example, soon after a person has had **unprotected sex** with a partner infected with HIV.

Detecting HIV at the earliest stage of infection lets people take steps right away to prevent **transmission of HIV**. (See the [Preventing Transmission of HIV](#) fact sheet.) This is important because immediately after infection the amount of HIV in the body is very high, increasing the risk of transmission of HIV. Starting treatment at this earliest stage of infection also can be considered.

### What does it mean to test HIV positive?

A diagnosis of HIV is made on the basis of positive results from **two** HIV tests. The first test can be either an HIV antibody test (using blood, urine, or fluids from the mouth) or a plasma HIV RNA test (using blood). The second test

(always using blood) is a different type of antibody test called a **Western blot** test. A positive Western blot test confirms that a person has HIV.

### How long does it take to get HIV test results?

Results of the first antibody test are generally available within a few days. (**Rapid HIV antibody tests** can produce results within an hour.) Results of the plasma HIV RNA test and Western blot are available in a few days to a few weeks.

### If I test HIV positive now, will I always test HIV positive?

Yes. There's no cure for HIV at this time. Because you will always be infected with the virus, you will always test HIV positive. But treatment with anti-HIV medications can help you live a longer, healthier life.

#### Terms Used in This Fact Sheet:

**HIV antibody test:** An HIV test that checks for HIV antibodies in a person's blood, urine, or fluids from the mouth. HIV antibodies are a type of protein the body produces in response to HIV infection.

**Mother-to-child transmission of HIV:** The passing of HIV from a woman infected with HIV to her baby during pregnancy, during labor and delivery, or by breastfeeding.

**Plasma HIV RNA test (viral load test):** A test that measures the amount of HIV in the blood. This test is used to detect recent HIV infection or to measure viral load at any stage of HIV infection.

**Rapid HIV antibody test:** An HIV antibody test that can detect HIV antibodies in blood or oral fluids in less than 30 minutes.

**Transmission of HIV:** The spread of HIV from a person infected with HIV to another person through the infected person's blood, semen, genital fluids, or breast milk.

**Unprotected sex:** Sex without using a condom.

**Viral load:** The amount of HIV in the blood. One of the goals of antiretroviral therapy is to reduce viral load.

**Western blot:** A type of antibody test used to confirm a positive HIV antibody or plasma HIV RNA test.

**Window period:** The time period between a person's infection with HIV and the appearance of detectable HIV antibodies.

## If a pregnant woman tests positive for HIV, will her baby be born with HIV?

In the United States and Europe, fewer than 2 babies in 100 born to mothers infected with HIV are infected with the virus. This is because anti-HIV medications given to women infected with HIV during pregnancy and delivery and to their babies after birth help prevent **mother-to-child transmission of HIV**. Another reason is that, in the United States and Europe, mothers infected with HIV do not breastfeed their babies. (For more information, see the [HIV and Pregnancy](#) fact sheet series.)

## Where can I find information on HIV testing in my state?

Many hospitals, medical clinics, and community organizations offer HIV testing. To find an HIV testing site near you, contact AIDSinfo for the number of your state AIDS hotline or visit <http://www.hivtest.org/>. You can also find information on testing locations on your state health department website.

## For more information:

Contact an AIDSinfo health information specialist at 1–800–448–0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.



## Seeing an HIV Health Care Provider

### I just tested HIV positive. What should I look for in a health care provider?

Look for a health care provider who has experience treating HIV and AIDS. You may want to see a specialist in HIV.

You need a health care provider with whom you feel comfortable. You will be working closely with your health care provider to make many decisions regarding your treatment.

### What can I expect at my first health care provider visit?

Your health care provider will ask you about your health and lifestyle, do a physical exam, and order blood tests. Your health care provider will also discuss what it means to have HIV and how it can affect your life. Your first visit is a good time to ask your health care provider questions.

### What questions should I ask my health care provider?

Ask your health care provider about:

- The benefits and risks of HIV treatment
- How HIV treatment can affect your lifestyle
- Lab tests used to monitor HIV infection
- How to avoid getting other infections
- How to avoid spreading HIV to another person

Write down your questions so you remember them when you visit your health care provider.

### What tests will my health care provider order?

You will have three very important blood tests at your first medical appointment: a **CD4 count**, a **viral load test**, and **drug-resistance testing**.

- A **CD4 count** is the number of CD4 cells in a sample of blood. CD4 cells are infection-fighting cells of the body's immune system. HIV destroys CD4 cells, making it hard for the body to fight off infections. A CD4 count measures how well the immune system is working. A goal of HIV treatment is to prevent HIV from destroying CD4 cells.
- A **viral load test** measures the amount of HIV in a sample of blood. The test indicates how much virus is in the blood (viral load). A goal of HIV treatment is to keep a person's viral load so low that the virus can't be detected by a viral load test.

- **Drug-resistance testing** identifies which, if any, anti-HIV medications will not be effective against a person's strain of HIV.

Your health care provider may also order other tests, such as a blood cell count, kidney and liver function tests, and tests for **sexually transmitted diseases (STDs)** and other diseases.

### When will I begin HIV treatment?

Starting HIV treatment is a big step. When to begin treatment depends on your health, your test results, and your readiness to take a combination of anti-HIV medications (a **regimen**) every day. Once you begin taking anti-HIV medications, you will probably need to take them for the rest of your life.

Your health care provider will help you decide if you are ready to start treatment. (See the [When to Start Anti-HIV Medications](#) fact sheet.) Once you start treatment, your health care provider will help you find ways to stick to your treatment regimen. (See the [Treatment Adherence](#) and [Following an HIV Treatment Regimen](#) fact sheets.)

### What happens if I don't start treatment right away?

If you don't start treatment right away, you should have a CD4 count and viral load test once every 3 to 6 months. Your health care provider will use the test results to monitor your infection and help you decide when to start treatment.

#### Terms Used in This Fact Sheet:

**CD4 count:** The number of CD4 cells in a sample of blood. A CD4 count measures how well the immune system is working.

**Drug-resistance testing:** A blood test to identify which, if any, anti-HIV medications will not be effective against a person's specific strain of HIV. Drug-resistance testing is done using a sample of blood.

**Regimen:** A combination of three or more anti-HIV medications from at least two different drug classes.

**Sexually transmitted diseases (STDs):** Infections that are usually passed during sex.

**Viral load:** The amount of HIV in the blood. One of the goals of antiretroviral therapy is to reduce viral load.

**For more information:**

Contact an AIDSinfo health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.





## When to Start Anti-HIV Medications

### I just tested HIV positive. When will I start treatment?

**Antiretroviral therapy (ART)** is recommended for all people infected with HIV. ART involves taking a combination of anti-HIV medications (a **regimen**) every day. ART is a lifelong treatment.

When to start anti-HIV medications (also called **antiretrovirals**) is a decision you will make with your health care provider. You and your health care provider will consider the following factors:

- How well your immune system is working (**CD4 count**)
- The amount of HIV in your blood (**viral load**)
- Whether you have an HIV-related illness or AIDS
- Whether you're pregnant
- Your ability and willingness to commit to lifelong treatment

### Can anti-HIV medications really help?

Yes. Anti-HIV medications can't cure HIV, but treatment can improve your quality of life and help you live longer.

HIV attacks and destroys the infection-fighting CD4 cells of the body's immune system. Loss of CD4 cells makes it hard for the body to fight infection. Anti-HIV medications can prevent HIV from multiplying. This reduces the amount of HIV in the body, giving the immune system a chance to recover and produce more infection-fighting CD4 cells. Once a person starts taking anti-HIV medications, an increase in CD4 cells is a sign that the immune system is recovering.

### How long does it take for treatment to work?

Once you start treatment—and take your anti-HIV medications exactly as directed—it's possible to have an **undetectable viral load** within 3 to 6 months. An undetectable viral load means that the level of HIV in your blood is too low to be detected by a viral load test. You aren't cured. There is still some HIV in your body. But an undetectable viral load indicates that your anti-HIV medications are working effectively to keep you healthier and reduce your risk of transmitting HIV.

#### Terms Used in This Fact Sheet:

**Antiretroviral:** A medication that prevents a retrovirus, such as HIV, from making copies of itself. Anti-HIV medications are also called antiretrovirals.

**Antiretroviral therapy (ART):** The recommended treatment for HIV. ART involves taking a combination of three or more anti-HIV medications from at least two different drug classes every day to control the virus.

**CD4 count:** The number of CD4 cells in a sample of blood. A CD4 count measures how well the immune system is working.

**Drug class:** A group of medications that work in the same way.

**Regimen:** A combination of three or more anti-HIV medications from at least two different drug classes.

**Undetectable viral load:** When the amount of HIV in a person's blood is too low to be detected with a viral load test.

**Viral load:** The amount of HIV in the blood. One of the goals of antiretroviral therapy is to reduce viral load.

### What treatment is right for me?

The U.S. Department of Health and Human Services (HHS) provides guidelines on using anti-HIV medications to treat HIV infection. The HHS guidelines recommend starting treatment with a regimen of three or more anti-HIV medications from at least two different **drug classes**. (See the [FDA-Approved Anti-HIV Medications](#) fact sheet.) The HHS guidelines list preferred ART regimens. (See the [Recommended HIV Treatment Regimens](#) fact sheet.) Because people's needs vary, the preferred regimens may not be right for everyone. You and your health care provider will consider your individual needs to select the most effective regimen for you.

### For more information:

Contact an AIDSinfo health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.

## Recommended HIV Treatment Regimens

### What is the treatment for HIV?

**Antiretroviral therapy (ART)** is the recommended treatment for HIV. ART involves taking a combination of anti-HIV medications (a **regimen**) every day. Anti-HIV medications (also called **antiretrovirals**) are grouped into six **drug classes** according to how they fight HIV. The six classes are non-nucleoside reverse transcriptase inhibitors (NNRTIs), nucleoside reverse transcriptase inhibitors (NRTIs), protease inhibitors (PIs), fusion inhibitors, CCR5 antagonists, and integrase inhibitors.

Recommended HIV treatment regimens include three or more anti-HIV medications from at least two different drug classes. Taking a combination of anti-HIV medications from different classes is the most effective way to control the virus. Some anti-HIV medications are available in combination (two or more medications in one pill).

Anti-HIV medications are approved by the U.S. Food and Drug Administration (FDA). See the [FDA-Approved Anti-HIV Medications](#) fact sheet for a complete list of medications used in HIV treatment regimens in the United States.

### How will I know which anti-HIV medications to take?

The best combination of anti-HIV medications for you depends on your individual needs. Factors that you and your health care provider will consider when selecting your HIV regimen include:

- Other diseases or conditions you may have
- Possible side effects of anti-HIV medications
- The risk of interactions between anti-HIV medications and other medications you take
- Results of **drug-resistance testing** and other tests
- Convenience of the regimen. (For example, a regimen that involves taking only one pill a day is convenient to follow.)
- Any personal issues that can make following a regimen difficult (such as depression or alcohol or drug abuse)

### What are the recommended regimens for people taking anti-HIV medications for the first time?

After considering your individual needs, you and your health care provider may select one of the following regimens

recommended for people taking anti-HIV medications for the first time:

- **Atripla** (a combination of three anti-HIV medications in one pill)
- **Reyataz + Norvir + Truvada** (Truvada is a combination of two anti-HIV medications in one pill.)
- **Prezista + Norvir + Truvada**
- **Isentress + Truvada**

#### Terms Used in This Fact Sheet:

**Antiretroviral:** A medication that prevents a retrovirus, such as HIV, from making copies of itself. Anti-HIV medications are also called antiretrovirals.

**Antiretroviral therapy (ART):** The recommended treatment for HIV. ART involves taking a combination of three or more anti-HIV medications from at least two different drug classes every day to control the virus.

**Atripla:** A combination of three anti-HIV medications in one pill—Sustiva (also called efavirenz or EFV), Emtriva (also called emtricitabine or FTC), and Viread (also called tenofovir or TDF).

**Drug class:** A group of medications that work in the same way.

**Drug-resistance testing:** Testing to identify which anti-HIV medications will or will not be effective against a person's specific strain of HIV. Drug-resistance testing is done using a sample of blood.

**Isentress:** An anti-HIV medication in the integrase inhibitor class. Isentress is also called raltegravir or RAL.

**Norvir:** An anti-HIV medication in the protease inhibitor (PI) class. Norvir is also called ritonavir or RTV.

**Prezista:** An anti-HIV medication in the protease inhibitor (PI) class. Prezista is also called darunavir or DRV.

**Regimen:** A combination of three or more anti-HIV medications from at least two different drug classes.

**Reyataz:** An anti-HIV medication in the protease inhibitor (PI) class. Reyataz is also called atazanavir or ATV.

**Sustiva:** An anti-HIV medication in the non-nucleoside reverse transcriptase inhibitor (NNRTI) class. Sustiva is also called efavirenz or EFV.

**Truvada:** Two anti-HIV medications from the nucleoside reverse transcriptase (NRTI) class—Emtriva and Viread—combined in a single pill. Emtriva is also called emtricitabine or FTC. Viread is also called tenofovir or TDF.

Women who are planning on becoming pregnant or are in the first trimester of pregnancy should not use Atripla or **Sustiva**. (Sustiva, which is one of the medications in Atripla, may cause birth defects that develop during the first few months of pregnancy.) If you are pregnant or expect to become pregnant soon, talk to your health care provider about the benefits and risks of taking anti-HIV medications. (See the [HIV and Pregnancy](#) fact sheet series for information on HIV treatment regimens for pregnant women.)

Because individual needs vary, these recommended HIV treatment regimens may not be right for everyone. If none of the preferred regimens is right for you, your health care provider will help you select an alternative regimen based on your needs.

### **Will I have side effects from the anti-HIV medications in my regimen?**

Anti-HIV medications can cause side effects. Side effects vary depending on the anti-HIV medication. And people taking the same medication may not have the same side effects. Before starting treatment, discuss possible side effects with your health care provider or pharmacist.

Most side effects from anti-HIV medications are manageable. However, side effects that become unbearable or life threatening call for a change in medications. Side effects that may seem minor, such as fever, nausea, fatigue, or rash, can indicate serious problems. Once you start treatment, always discuss any side effects from your anti-HIV medications with your health care provider.

Interactions between anti-HIV medications and other medications can increase the risk of side effects. Drug interactions can also reduce the effectiveness of anti-HIV medications. (Anti-HIV medications can also have the same effect on other medications.) Always tell your health care provider about other medications you take, including when you switch or stop taking a medication.

### **For more information:**

Contact an *AIDSinfo* health information specialist at 1–800–448–0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.

## FDA-Approved Anti-HIV Medications

Antiretroviral therapy (ART) is the recommended treatment for HIV infection. ART involves taking a combination of anti-HIV medications (a regimen) daily. A regimen contains three or more anti-HIV medications from at least two different drug classes. Anti-HIV medications prevent HIV from multiplying in the body, which helps people infected with HIV live longer, healthier lives. ART may reduce the risk of transmission of HIV but anti-HIV medications can't cure HIV/AIDS.

The following table lists anti-HIV medications approved by the U.S. Food and Drug Administration (FDA) for treatment of HIV in the United States. The medications are presented by drug class and identified by generic name/acronym and brand name.

Drug Class	Generic Name (Acronym)	Brand Name	Manufacturer	FDA Approval Date
<b>Non-nucleoside Reverse Transcriptase Inhibitors (NNRTIs)</b>				
<i>NNRTIs bind to and alter reverse transcriptase, an enzyme HIV needs to make copies of itself.</i>	Delavirdine (DLV)	Rescriptor	Pfizer	April 4, 1997
	Efavirenz (EFV)	Sustiva	Bristol-Myers Squibb	Sept. 17, 1998
	Etravirine (ETR)	Intelence	Tibotec	Jan. 18, 2008
	Nevirapine (NVP)	Viramune	Boehringer Ingelheim	June 21, 1996
	Rilpivirine (RPV)	Edurant	Janssen Pharmaceuticals, Inc.	May 20, 2011
<b>Nucleoside Reverse Transcriptase Inhibitors (NRTIs)</b>				
<i>NRTIs block reverse transcriptase, an enzyme HIV needs to make copies of itself.</i>	Abacavir (ABC)	Ziagen	GlaxoSmithKline	Dec. 17, 1998
	Didanosine (ddI)	Videx Videx EC (enteric-coated)	Bristol-Myers Squibb Bristol-Myers Squibb	Oct. 9, 1991 Oct. 31, 2000
	Emtricitabine (FTC)	Emtriva	Gilead Sciences	July 2, 2003
	Lamivudine (3TC)	Epivir	GlaxoSmithKline	Nov. 17, 1995
	Stavudine (d4T)	Zerit	Bristol-Myers Squibb	June 24, 1994
	Tenofovir DF (TDF)	Viread	Gilead Sciences	Oct. 26, 2001
	Zidovudine (ZDV, AZT)	Retrovir	GlaxoSmithKline	March 19, 1987
<b>Protease Inhibitors (PIs)</b>				
<i>PIs block HIV protease, an enzyme HIV needs to make copies of itself.</i>	Atazanavir (ATV)	Reyataz	Bristol-Myers Squibb	June 20, 2003
	Darunavir (DRV)	Prezista	Janssen Pharmaceuticals, Inc.	June 23, 2006
	Fosamprenavir (FPV)	Lexiva	GlaxoSmithKline	Oct. 20, 2003
	Indinavir (IDV)	Crixivan	Merck	March 13, 1996
	Nelfinavir (NFV)	Viracept	Agouron Pharmaceuticals	March 14, 1997

Drug Class	Generic Name (Acronym)	Brand Name	Manufacturer	FDA Approval Date
<b>Protease Inhibitors (PIs), continued</b>				
<i>PIs block HIV protease, an enzyme HIV needs to make copies of itself.</i>	Ritonavir (RTV)	Norvir	Abbott Laboratories	March 1, 1996
	Saquinavir (SQV)	Invirase	Hoffmann-La Roche	Dec. 6, 1995
	Tipranavir (TPV)	Aptivus	Boehringer Ingelheim	June 20, 2005
<b>Fusion Inhibitors</b>				
<i>Fusion inhibitors block HIV from entering the CD4 cells of the immune system.</i>	Enfuvirtide (T-20)	Fuzeon	Hoffmann-La Roche, Trimeris	March 13, 2003
<b>CCR5 Antagonists</b>				
<i>CCR5 entry inhibitors block CCR5, a protein on the CD4 cells that HIV needs to enter the cells.</i>	Maraviroc (MVC)	Selzentry	Pfizer	Aug. 6, 2007
<b>Integrase Inhibitors</b>				
<i>Integrase inhibitors block HIV integrase, an enzyme HIV needs to make copies of itself.</i>	Raltegravir (RAL)	ISENTRESS	Merck	Oct. 12, 2007
<b>Fixed-Dose Combination</b>				
Fixed-dose combination tablets contain two or more anti-HIV medications from one or more drug classes.	Abacavir, Lamivudine	Epzicom	GlaxoSmithKline	Aug. 2, 2004
	Abacavir, Lamivudine, Zidovudine	Trizivir	GlaxoSmithKline	Nov. 14, 2000
	Efavirenz, Emtricitabine, Tenofovir DF	Atripla	Bristol-Myers Squibb, Gilead Sciences	July 12, 2006
	Emtricitabine, Rilpivirine, Tenofovir DF	Complera	Gilead Sciences	Aug. 10, 2011
	Emtricitabine, Tenofovir DF	Truvada	Gilead Sciences	Aug. 2, 2004
	Lamivudine, Zidovudine	Combivir	GlaxoSmithKline	Sept. 27, 1997
	Lopinavir, Ritonavir	Kaletra	Abbott Laboratories	Sept. 15, 2000



## Treatment Adherence

### What is treatment adherence?

**Treatment adherence** means following your treatment regimen closely every day—taking the correct dose of each anti-HIV medication at the correct time and exactly as prescribed. Adherence is very important for successful HIV treatment.

### Why is adherence important?

Adherence affects HIV treatment in two ways:

- Close adherence to an HIV treatment regimen allows anti-HIV medications to work effectively to reduce the amount of HIV in the body. Skipping medications, even occasionally, gives HIV the chance to multiply rapidly. Preventing the virus from multiplying is the best way to protect your health.
- Close adherence to an HIV treatment regimen also helps prevent **drug resistance**. Drug resistance develops when the virus mutates (changes form), becoming “resistant” to certain anti-HIV medications. One or more anti-HIV medications in a treatment regimen can become ineffective as a result of drug resistance.

Skipping medications makes it easier for drug resistance to develop. HIV can become resistant to the anti-HIV medications in a person’s current regimen or to other, similar anti-HIV medications not yet taken, limiting options for successful HIV treatment. And drug-resistant strains of HIV can be transmitted to others, too.

There are many different anti-HIV medications and treatment regimens, but studies show that **a person’s first regimen offers the best chance for long-term treatment success**. And good adherence from the start is key to successful treatment.

### Why is treatment adherence sometimes difficult?

Adhering to an HIV treatment regimen can be difficult for several reasons. Some treatment regimens involve taking several pills every day—with or without food, or before or after other medications. Other factors that can make treatment adherence difficult include:

- Difficulty taking medications (such as trouble swallowing pills)
- Side effects from medications (for example, nausea or diarrhea)

- A busy schedule, shift work, or travel away from home that makes it easy to forget to take pills
- Being sick or depressed
- Alcohol or drug abuse

### What can I do to adhere to my HIV treatment regimen?

Before you start treatment, be certain you’re committed to taking anti-HIV medications every day as directed. Talk to your health care provider about any issues that can make adherence difficult, including:

- Possible side effects from the anti-HIV medications in your regimen
- How other medications you take may interact with your anti-HIV medications
- Your schedule at home and at work
- Any personal issues such as depression or alcohol or drug abuse
- Lack of health insurance to pay for anti-HIV medications

Understanding issues that can make adherence difficult will help you and your health care provider select the best regimen for you. Some people find that adhering to an HIV treatment regimen becomes more difficult over time. So, every time you see your health care provider, make it a point to talk about adherence. (See the [Following an HIV Treatment Regimen](#) fact sheet for tips on adherence.)

### For more information:

Contact an AIDSinfo health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.

#### Terms Used in This Fact Sheet:

**Drug resistance:** When HIV mutates (changes form), causing one or more anti-HIV medications to be ineffective.

**Regimen:** A combination of three or more anti-HIV medications from at least two different drug classes.

**Treatment adherence:** Closely following an HIV treatment regimen—taking the correct dose of each anti-HIV medication at the correct time and exactly as prescribed.

## Following an HIV Treatment Regimen

### How can I prepare for adherence before I start HIV treatment?

Preparing for adherence before you start taking anti-HIV medications is the first step to treatment success. Planning ahead will help you follow your treatment **regimen** once you start treatment.

Begin by talking to your health care provider. Make sure you understand why you're starting HIV treatment and why **treatment adherence** is important. Discuss these important details about your treatment regimen:

- Each anti-HIV medication in your regimen
- The dose (amount) of each anti-HIV medication in your regimen.
- How many pills in each dose
- When to take each medication
- How to take each medication—with or without food
- Possible side effects from each medication, including serious side effects
- How to store your medications

Talk to your health care provider about other medications you take and their possible side effects. Your health care provider will tell you about potential interactions between the anti-HIV medications in your regimen and the other medications you take.

Tell your health care provider if you have any personal issues, such as depression or alcohol or drug abuse, that can make adherence difficult. If needed, your health care provider can recommend resources to help you address these issues before you start treatment.

### How can I maintain adherence after I start treatment?

Consider one or more of the following strategies to help you adhere to your regimen:

- Use a 7-day pill box. Once a week, fill the pill box with your medications for the entire week.
- Take your medications at the same time every day.
- Use a timer, an alarm clock, or your cell phone alarm to remind you to take your medications.
- Ask your family members, friends, or coworkers to remind you to take your medications.
- Keep your medications nearby. Keep a backup supply of

medications at work or in your purse or briefcase.

- Plan ahead for changes in your daily routine, including weekends and holidays. If you're going away, pack enough medications to last the entire trip.
- Use a medication diary to stay on track. Write down the name of each medication; include the dose, number of pills to take, and when to take them. Check off each medication as you take it. Reviewing your diary will help you identify the times you're most likely to skip medications.
- Keep all your medical appointments. Write down the date and time of health care provider visits on your calendar or daily schedule. If you run low on medications before your next visit, call your health care provider to renew your prescriptions.
- Get additional tips on adherence by joining a support group for people living with HIV.

### What should I do if I forget to take my medications?

Unless your health care provider tells you otherwise, take a medication you missed as soon as you realize you skipped it. But if it's almost time for the next dose of the medication, don't take the missed dose and just continue on your regular medication schedule. Don't take a double dose of a medication to make up for a missed dose.

### What should I do if I have problems adhering to my treatment regimen?

Tell your health care provider that you're having difficulty following your regimen. Together you can identify the reasons why you're skipping medications.

Tell your health care provider about any side effects from the medications in your regimen. Side effects are a major reason treatment adherence can be difficult. A regimen that involves taking many pills at many times during the day can also make adherence difficult.

#### Terms Used in This Fact Sheet:

**Regimen:** A combination of three or more anti-HIV medications from at least two different drug classes.

**Treatment adherence:** Closely following an HIV treatment regimen—taking the correct dose of each anti-HIV medication at the correct time and exactly as prescribed.

Based on why you're having problems with adherence, your health care provider may adjust or change your regimen. (See the [Changing an HIV Treatment Regimen](#) fact sheet.)

**For more information:**

Contact an AIDSinfo health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.





## Is My Treatment Regimen Working?

### How will I know if my HIV treatment regimen is working?

Your health care provider will use two important blood tests to monitor your HIV treatment: **CD4 count** and **viral load** test. The results of the tests will help your health care provider determine if the anti-HIV medications in your treatment **regimen** are working.

### What is a CD4 count?

HIV attacks the immune system, destroying the system's infection-fighting CD4 cells. Keeping the immune system healthy is an important goal of HIV treatment.

The CD4 count measures the number of CD4 cells in a sample of blood. The CD4 count of a healthy person ranges from 500 to 1,200 cells/mm<sup>3</sup>. An HIV-infected person with a CD4 count of less than 200 cells/mm<sup>3</sup> has AIDS.

Because a falling CD4 count is a sign that HIV is damaging the immune system, the test is used to monitor HIV infection. Once treatment is started, the CD4 count is also used to monitor the effectiveness of anti-HIV medications.

Once you start treatment, you should have a CD4 count once every 3 to 4 months. An increasing CD4 count is a sign that the immune system is recovering. If your regimen is working well, you need a CD4 count only once every 6 to 12 months.

### What is a viral load test?

Preventing HIV from multiplying is another important goal of HIV treatment. The viral load test measures the amount of HIV in the blood. It's the best measure of how well anti-HIV medications are controlling the virus.

The best sign that treatment is working is reaching and maintaining an **undetectable viral load**. An undetectable viral load doesn't mean that you're cured. It means that the amount of HIV in your blood is too low to be detected by the viral load test.

Once you start treatment, you should have a viral load test within 2 to 8 weeks and then once every 4 to 8 weeks until your viral load is undetectable. You need the test done only every 3 to 4 months once your viral load is undetectable. If you have an undetectable viral load for more than 2 or 3 years, your health care provider may recommend viral load testing once every 6 months.

### What causes treatment to fail?

HIV treatment can fail if anti-HIV medications are unable to control the virus or protect the health of the immune system.

Sometimes treatment fails because of things you can't control, such as unmanageable side effects from anti-HIV medications, interactions between anti-HIV medications and other medications you take, or the body's poor absorption of anti-HIV medications.

Treatment can also fail because of **drug resistance**. Sometimes HIV changes form and becomes resistant to (not affected by) the medications in a regimen.

It may be necessary to change medications to deal with these problems.

### Can skipping medications cause treatment failure?

Poor **treatment adherence** is another reason HIV treatment can fail. Skipping medications allows HIV to multiply, increasing a person's viral load. To reach and maintain an undetectable viral load, it's important to closely follow your treatment regimen. Poor treatment adherence can also give HIV a chance to change form, leading to drug resistance.

Sometimes working with your health care provider to improve adherence can prevent treatment failure. For

#### Terms Used in This Fact Sheet:

**CD4 count:** The number of CD4 cells in a sample of blood. A CD4 count measures how well the immune system is working.

**Drug resistance:** When HIV mutates (changes form), causing one or more anti-HIV medications to be ineffective.

**Regimen:** A combination of three or more anti-HIV medications from at least two different drug classes.

**Treatment adherence:** Closely following an HIV treatment regimen—taking the correct dose of each anti-HIV medication at the correct time and exactly as prescribed.

**Undetectable viral load:** The amount of HIV in a person's blood is too low to be detected with a viral load test.

**Viral load:** The amount of HIV in the blood. One of the goals of antiretroviral therapy is to reduce viral load.

example, your health care provider can give you tips on how to manage medication side effects that make adherence difficult. Or your health care provider can simplify your regimen to make your medication schedule fit your busy lifestyle. (To learn more about treatment adherence, see the [Treatment Adherence](#) and [Following an HIV Treatment Regimen](#) fact sheets.)

### What happens if my treatment fails?

If your treatment is failing, it may be time to adjust or change your regimen. But before making any changes, your health care provider will consider:

- Any side effects you had from your anti-HIV medications
- How well your body absorbed the medications in your regimen

- Your drug-resistance testing results
- How closely you followed your treatment regimen

All of this information will help you and your health care provider select a new, more effective regimen. (See the [Changing an HIV Treatment Regimen](#) fact sheet.)

### For more information:

Contact an *AIDSinfo* health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.

## Changing an HIV Treatment Regimen

### Will my HIV treatment regimen ever change?

At some point, you may need to adjust or change your **regimen**. But before making any changes, it's important to understand why.

### What are possible reasons for changing an HIV treatment regimen?

There are several reasons why a person may switch to another HIV regimen:

- **Side effects from anti-HIV medications**  
Unpleasant side effects, such as fatigue, nausea, and diarrhea, can make **treatment adherence** difficult. Side effects that become unbearable or pose a serious threat to health call for a change in regimen.
- **Poor absorption of anti-HIV medications**  
To work effectively, anti-HIV medications must be absorbed by the body.
- **Drug interactions**  
**Drug interactions** between anti-HIV medications in a regimen or between anti-HIV medications and other medications a person is taking can increase the risk of side effects. Drug interactions can also reduce the effectiveness of anti-HIV medications. (Anti-HIV medications can also have the same effect on other medications.)
- **Drug resistance**  
**Drug resistance** occurs when HIV mutates (changes form), causing one or more medications in a regimen to be ineffective.
- **Poor treatment adherence**  
Skipping medications gives HIV the chance to multiply, increasing a person's **viral load**. Poor adherence also increases the risk of drug resistance.

### What are important things to consider when selecting a new treatment regimen?

If you and your health care provider decide it's time to switch your treatment regimen, you will have many things to consider. For example, together you will review:

- your medication history
- any side effects from the anti-HIV medications you currently take or have taken in the past
- results of **drug-resistance testing**

In general, a new treatment regimen should include two or

more medications from two or more **drug classes**. If you are switching regimens, your new regimen may include anti-HIV medications that you have never used before.

If you have already taken many of the FDA-approved anti-HIV medications, your health care provider may recommend a new medication only available through a research study (**clinical trial**). To learn about participating in a research study, ask your health care provider or visit the Clinical Trials section of the *AIDSinfo* website at <http://aidsinfo.nih.gov/clinicaltrials>.

### How can I give my new regimen the best chance of success?

Before starting your new regimen, make a commitment to keep your medical appointments and take your anti-HIV medications exactly as prescribed. Talk to your health care provider about steps you can take to overcome any lifestyle or personal issues that can make adherence difficult.

(See the [Treatment Adherence](#) and [Following an HIV Treatment Regimen](#) fact sheets.)

#### Terms Used in This Fact Sheet:

**Clinical trial:** A type of research study that tests how well medical treatments work in people.

**Drug class:** A group of medications that work in the same way.

**Drug interaction:** A change in how a drug works when taken with another drug (drug-drug interaction) or with a specific food (food-drug interaction).

**Drug resistance:** When HIV mutates (changes form), causing one or more anti-HIV medications to be ineffective.

**Drug-resistance testing:** A blood test to identify which, if any, anti-HIV medications will not be effective against a person's specific strain of HIV. Drug-resistance testing is done using a sample of blood.

**Regimen:** A combination of three or more anti-HIV medications from at least two different drug classes.

**Treatment adherence:** Closely following an HIV treatment regimen—taking the correct dose of each anti-HIV medication at the correct time and exactly as prescribed.

**Viral load:** The amount of HIV in the blood. One of the goals of antiretroviral therapy is to reduce viral load.

Be sure to ask your health care provider about possible side effects from your new anti-HIV medications. Also discuss potential drug interactions between the medications in your regimen and other medications, vitamins, nutritional supplements, and herbal products that you take or plan to take.

**For more information:**

Contact an *AIDSinfo* health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.

## HIV Coinfections

### What is a coinfection?

**Coinfection** means infection with more than one disease at the same time. Some coinfections commonly seen in people infected with HIV include:

- HIV/**hepatitis B virus (HBV)** coinfection
- HIV/**hepatitis C virus (HCV)** coinfection
- HIV/**tuberculosis (TB)** coinfection

People infected with HIV should be tested for HBV, HCV, and TB.

### What are HBV and HCV?

HBV and HCV are two different viruses that both cause liver disease. They are also among the most common causes of liver cancer. Some of the ways HBV, HCV, and HIV are spread are similar.

HBV is spread through the blood, semen, or other body fluid of a person infected with HBV. Having **unprotected sex** or sharing drug injection equipment (such as needles or syringes) with a person infected with HBV are the main ways people get HBV. (To prevent HBV infection, people infected with HIV receive the HBV vaccination.)

HCV is spread through the blood of a person infected with HCV. Sharing drug injection equipment with a person infected with HCV is the main way people get HCV, but HCV can also be transmitted during unprotected sex. (Before widespread screening of the blood supply began in 1992, HCV was also commonly spread through blood transfusions and organ transplants.)

Having unprotected sex or sharing drug needles are also ways people get HIV. That is why some people become coinfecting with HIV and HBV or HCV (or both) at the same time.

### What is TB?

TB is a disease caused by germs that spread through the air when a person with active TB coughs, sneezes, or talks. TB usually affects the lungs.

There are two forms of TB: **latent TB infection** and **TB disease**. Latent TB infection is the inactive form of TB. The TB germs in the body are “sleeping” and don’t make the person sick. A person with latent TB infection can’t spread TB to others.

Without treatment, latent TB infection can advance to TB

disease, especially in people with weakened immune systems. The TB germs in the body multiply and become active, making the person sick. A person with TB disease of the lungs can spread TB to others.

Because HIV weakens the immune system, latent TB infection is more likely to advance to TB disease in a person infected with HIV. In a person infected with HIV, TB disease is considered an **AIDS-defining condition**, and TB treatment should be started immediately.

### Are coinfections more serious in people infected with HIV?

Yes. Coinfections can become serious more rapidly in people infected with HIV than in people who are not infected with the virus.

HBV and HCV both lead to liver damage more quickly in people infected with HIV. People co-infected with HBV or

#### Terms Used in This Fact Sheet:

**AIDS-defining condition:** Any of several illnesses that can lead to a diagnosis of AIDS in a person infected with HIV. AIDS is the most advanced stage of HIV infection.

**Coinfection:** Infection with more than one disease at the same time. Some people infected with HIV are coinfecting with hepatitis B virus (HBV), hepatitis C virus (HCV), or tuberculosis (TB).

**Hepatitis B virus (HBV):** The virus that causes a disease of the liver (hepatitis B).

**Hepatitis C virus (HCV):** The virus that causes a disease of the liver (hepatitis C).

**Latent tuberculosis (TB) infection:** The inactive form of TB, which doesn’t make a person sick and can’t be spread to other people.

**Tuberculosis (TB) disease:** The active form of TB, which makes a person sick and can be spread to other people if the infection involves the lungs. In a person infected with HIV, TB disease is considered an AIDS-defining condition.

**Tuberculosis (TB):** A disease caused by germs that spread through the air when a person with active TB coughs, sneezes, or talks. TB usually affects the lungs.

**Unprotected sex:** Sex without using a condom.

HCV also have a higher risk of developing liver damage from anti-HIV medications.

TB disease is more likely to spread beyond the lungs in people infected with HIV than in people who do not have HIV.

### Can coinfections be treated?

Yes, but the effectiveness of treatment depends on the coinfection.

- TB treatment can cure TB disease or prevent latent TB infection from advancing to TB disease.
- There is no cure for HBV, but treatment can slow down HBV infection.
- Treatment for HCV is generally less effective than treatment for TB or HBV. However, research on new medications that are more effective against HCV is underway.

### Are HIV and coinfections treated at the same time?

Yes, but what medications to take and when to start them depend on the coinfection. Some anti-HIV medications are

effective against both HIV and HBV. Treatment for HCV or TB involves taking other medications in addition to anti-HIV medications.

Health care providers closely watch people receiving treatment for coinfections for any side effects from anti-HIV medications or medications used to treat coinfections. They also watch for drug interactions between the medications. Changing medications can be helpful to avoid side effects or drug interactions.

Talk to your health care provider if you have questions about HIV and coinfections.

### For more information:

Contact an *AIDSinfo* health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.



## Preventing Transmission of HIV

### How is HIV transmitted?

HIV is transmitted (spread) through the blood, semen, genital fluids, or breast milk of a person infected with HIV. The spread of the virus is called **transmission of HIV**.

Having **unprotected sex** or sharing drug injection equipment (such as needles and syringes) with a person infected with HIV are the most common ways HIV is transmitted.

Having a **sexually transmitted disease (STD)** can increase a person's risk of becoming infected with HIV during sex. The risk of spreading HIV during sex is also more likely if the partner infected with HIV also has another STD.

Women infected with HIV can transmit the virus to their babies during pregnancy or childbirth or by breastfeeding. If you are a woman infected with HIV, talk to your health care provider about ways to prevent pregnancy. If you are pregnant or plan to become pregnant, ask your health care provider how you can protect your baby from HIV. (See the [HIV and Pregnancy](#) fact sheets.)

### I am taking anti-HIV medications and my viral load is undetectable. Can I still infect another person with HIV?

Your anti-HIV medications are doing a good job of controlling your infection. The amount of HIV in your blood is so low that a **viral load** test can't detect the virus. But having an **undetectable viral load** doesn't mean you're cured. You still have HIV. Although having an undetectable viral load greatly reduces the risk of HIV transmission, you can still infect another person with the virus.

### How can I prevent transmitting HIV?

To prevent infecting another person with HIV:

- Use a condom every time you have sex.
- If you inject drugs, don't share your needles or syringes.
- Don't share your razor, toothbrush, or other items that may have your blood on them.
- Take your anti-HIV medications according to your health care provider's directions.
- If you are a mother infected with HIV, don't breastfeed your baby.

Talk to your health care provider about how HIV is transmitted and ways to prevent spreading the virus. At each visit, discuss any high-risk behaviors (such as having unprotected sex or sharing drug injection equipment). Ask

your health provider about testing for other STDs—for you and your partner.

Talking about high-risk behaviors can be difficult. But it's important to be honest with your health care provider about any high-risk activities. Your health care provider can help you take steps to reduce your chances of transmitting HIV to another person.

### Can I put my HIV-infected partner at risk?

Even if your partner is also infected with HIV, it's important to use condoms and not share drug injection equipment. You and your partner may have different strains of the virus. Your partner's HIV could act differently in your body or cause the anti-HIV medications you take to be less effective. And your strain of HIV could have the same effects on your partner.

### Where can I find more information about HIV prevention?

The Centers for Disease Control and Prevention (CDC) National Prevention Information Network (NPIN) provides information about the prevention of HIV infection, other STDs, and **tuberculosis (TB)**.

If you have questions about HIV transmission, call CDC-INFO at 1-800-232-4636 or visit <http://www.cdc.gov/hiv/>.

### For more information:

Contact an *AIDSinfo* health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.

#### Terms Used in This Fact Sheet:

**Sexually transmitted diseases (STDs):** Infections that are usually passed during sex. HIV is an example of an STD.

**Transmission of HIV:** The spread of HIV from a person infected with HIV to another person through the infected person's blood, semen, genital fluids, or breast milk.

**Tuberculosis (TB):** A disease caused by germs that spread through the air when a person with active TB coughs, sneezes, or talks. TB usually affects the lungs.

**Undetectable viral load:** When the amount of HIV in a person's blood is too low to be detected with a viral load test.

**Unprotected sex:** Sex without using a condom.

**Viral load:** The amount of HIV in the blood. One of the goals of antiretroviral therapy is to reduce viral load.