Sulfamethoxazole / Trimethoprim

sul-fa-meth-OX-a-zole, trye-METH-oh-prim
Brand Name: Septra, Sulfatrim, Bactrim

Drug Class: Opportunistic Infection and Other Drugs



Sulfamethoxazole-trimethoprim, also known as Bactrim, Septra, or SMX-TMP, belongs to the class of medicines called antibacterials. It also belongs to the class of medicines called antiprotozoals. Antibacterials kill bacteria or stop bacteria from multiplying. Antiprotozoals help the body overcome infections caused by protozoa, which are microscopic organisms that can cause disease. When bacterial and protozoal infections occur in patients with weakened immune systems, including people with HIV or AIDS, they tend to be more serious and are called opportunistic infections (OIs).

HIV/AIDS-Related Uses

Sulfamethoxazole-trimethoprim was approved by the FDA on June 23, 1981, for the treatment and prevention of Pneumocystis carinii pneumonia (PCP), a common OI in people with HIV.

Dosage Form/Administration

Sulfamethoxazole-trimethoprim comes in oral liquid, tablet, and injection forms. The oral liquid and tablet forms should be taken with a full glass of water. The oral liquid should be shaken well before use.

Contraindications

Individuals should tell a doctor about any medical problems before taking this medicine.

Possible Side Effects

Along with its desired effects, sulfamethoxazole-trimethoprim can cause some unwanted effects. Serious side effects of this medicine include itching; skin rash; aching of joints and muscles; difficulty in swallowing; pale skin; redness; blistering, peeling, or loosening of skin; sore throat and fever; unusual bleeding or bruising; unusual tiredness or weakness; yellow eyes or skin; or increased sensitivity of skin to sunlight. Individuals should tell a doctor if they have any of these side effects.

Other side effects may not be serious and may lessen or disappear with continued use of the medicine. Less serious side effects of this medicine include diarrhea, dizziness, headache, loss of appetite, nausea or vomiting, and tiredness. Individuals should tell a doctor if these side effects continue or are bothersome.

Drug and Food Interactions

A doctor should be notified of any other medications being taken, including prescription, nonprescription (over-the-counter), or herbal medications.

Clinical Trials

For information on clinical trials that involve Sulfamethoxazole / Trimethoprim, visit the ClinicalTrials.gov web site at http://www.clinicaltrials.gov. In the Search box, enter: Sulfamethoxazole / Trimethoprim AND HIV Infections.

Manufacturer Information

Septra

King Pharmaceuticals, Inc. 501 Fifth Street Bristol, TN 37620 (888) 840-5370

Sulfamethoxazole / Trimethoprim King Pharmaceuticals, Inc. 501 Fifth Street Bristol, TN 37620 (888) 840-5370

Bactrim

Women First Healthcare 5355 Mira Sorrento Place - Suite 700 San Diego, CA 92121 (858) 509-1171

Sulfatrim

Alpharma, Inc. One Executive Drive Fort Lee, NJ 07024 (201) 947-7774

Sulfamethoxazole / Trimethoprim



For More Information

Contact your doctor or an AIDSinfo Health Information Specialist:

- Via Phone: 1-800-448-0440 Monday Friday, 12:00 p.m. (Noon) 5:00 p.m. ET
- Via Live Help: http://aidsinfo.nih.gov/live_help Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET