



\* Each drug is recommended for use in combination with other antiretrovirals. Dose may be adjusted when taken in combination

Nucleoside and/or Nucleotide Reverse Transcriptase Inhibitor (NRTI, NtRTI)
Non-nucleoside Reverse Transcriptase Inhibitor (NNRTI)
Protease Inhibitor (PI)
Fusion Inhibitor

Thanks to Andrew Halbur, BSPHarm



**COMBIVIR**  
(lamivudine/zidovudine)

**ADULT SINGLE DOSE\*:**  
One tablet (150 mg lamivudine, 300 mg zidovudine), twice daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**EMTRIVA**  
(emtricitabine, FTC)

**ADULT SINGLE DOSE\*:**  
One 200 mg capsule daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**EPIVIR**  
(lamivudine, 3TC)

**ADULT SINGLE DOSE\*:**  
One 300 mg tablet, once daily or one 150 mg tablet, twice daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**EPIZCOM**  
(abacavir/lamivudine)

**ADULT SINGLE DOSE\*:**  
One tablet (600 mg abacavir, 300 mg lamivudine), once daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**HIVID**  
(zalcitabine, ddC)

**ADULT SINGLE DOSE\*:**  
One 0.75 mg tablet, three times a day

**FOOD AND LIQUID QUALIFICATIONS:** Take on an empty stomach



**RETROVIR**  
(zidovudine, AZT)

**ADULT SINGLE DOSE\*:**  
One 300 mg tablet, twice daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**TRIZIVIR**  
(abacavir/lamivudine/zidovudine)

**ADULT SINGLE DOSE\*:**  
One tablet (300 mg abacavir, 150 mg lamivudine, 300 mg zidovudine), twice daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**TRUVADA**  
(tenofovir/emtricitabine)

**ADULT SINGLE DOSE\*:**  
One tablet (300 mg tenofovir and 300 mg emtricitabine), once daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**VIDEX EC**  
(didanosine, ddI)

**ADULT SINGLE DOSE\*:**  
One 400 mg capsule daily with adjustments for weight

**FOOD AND LIQUID QUALIFICATIONS:** Take on an empty stomach, at least one hour before or 2 hours after food or drink, except water



**VIREAD**  
(tenofovir)

**ADULT SINGLE DOSE\*:**  
One 300 mg tablet daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**ZERIT**  
(stavudine, d4T)

**ADULT SINGLE DOSE\*:**  
One 40 mg capsule, twice daily with adjustments for weight

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**ZIAGEN**  
(abacavir sulfate)

**ADULT SINGLE DOSE\*:**  
Two 300 mg tablets once daily (or one 300 mg tablet, twice daily)

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**RESCRIPTOR**  
(delavirdine)

**ADULT SINGLE DOSE\*:**  
Four 100 mg tablets, three times a day, or two 200 mg tablets, three times a day

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions



**SUSTIVA**  
(efavirenz)

**ADULT SINGLE DOSE\*:**  
One 600 mg tablet daily

**FOOD AND LIQUID QUALIFICATIONS:** Take on an empty stomach, or may be taken with a light, low-fat snack



**VIRAMUNE**  
(nevirapine)

**ADULT SINGLE DOSE\*:**  
One 200 mg tablet daily for the first 14 days, then one 200 mg tablet, twice daily, or two 200 mg tablets once daily

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)

**PHOTO OF 50 MG CAPSULE NOT AVAILABLE**

**AGENERASE**  
(amprenavir)

**ADULT SINGLE DOSE\*:**  
1200 mg twice daily (150 mg capsule no longer available due to new formulation Lexiva; 50 mg capsule and liquid still available)

**FOOD AND LIQUID QUALIFICATIONS:** May be taken with or without food, but avoid a high fat meal



**APTIVUS**  
(tipranavir)

**ADULT SINGLE DOSE\*:**  
Two 250 mg capsules with two 100 mg capsules Norvir twice daily

**FOOD AND LIQUID QUALIFICATIONS:** Take with food



**CRIVAN**  
(indinavir)

**ADULT SINGLE DOSE\*:**  
Two 400 mg capsules, every 8 hours (almost always boosted with Norvir, see Drug Guide)

**FOOD AND LIQUID QUALIFICATIONS:** Take with no food or a low-fat snack (with food with Norvir); drink at least 1.5 liters of water daily



**FORTOVASE**  
(saquinavir soft-gel)

**ADULT SINGLE DOSE\*:**  
Discontinued early 2006. Six 200 mg capsules, three times a day (almost always boosted with Norvir, see Drug Guide)

**FOOD AND LIQUID QUALIFICATIONS:** Take with food, or within two hours after a meal



**INVIRASE**  
(saquinavir hard-gel)

**ADULT SINGLE DOSE\*:**  
Two 500 mg tablets with one 100 mg capsule Norvir twice a day. (Or five 200 mg hard-gel capsules and 100 mg Norvir, twice daily.) Saquinavir must be dosed with Norvir.

**FOOD AND LIQUID QUALIFICATIONS:** Take with food, or within two hours after a meal



**KALETRA**  
(lopinavir/ritonavir)

**ADULT SINGLE DOSE\*:**  
Two tablets (200 mg lopinavir/50 mg ritonavir) twice daily. (Or three capsules, 133.33 mg lopinavir/33.3 mg ritonavir, twice daily; to be phased out in early 2006.)

**FOOD AND LIQUID QUALIFICATIONS:** Take with food, if possible



**LEXIVA**  
(fos-amprenavir calcium)

**ADULT SINGLE DOSE\*:**  
Two 700 mg capsules plus 200 mg Norvir, once daily (if protease inhibitor experienced, 700 mg with 100 mg Norvir twice daily)

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)



**NORVIR**  
(ritonavir)

**ADULT SINGLE DOSE\*:**  
Used mostly for boosting, once or twice a day with another PI

**FOOD AND LIQUID QUALIFICATIONS:** Take with food, if possible



**REYATAZ**  
(atazanavir sulfate)

**ADULT SINGLE DOSE\*:**  
Two 200 mg capsules daily (if treatment experienced: 300 mg with 100 mg Norvir daily)

**FOOD AND LIQUID QUALIFICATIONS:** Take with food



**VIRACEPT**  
(nelfinavir)

**ADULT SINGLE DOSE\*:**  
Two 625 mg tablets, twice daily (or five 250 mg tablets, twice daily)

**FOOD AND LIQUID QUALIFICATIONS:** Take with a meal



**FUZEON**  
(T-20)

**ADULT SINGLE DOSE\*:**  
90 mg twice daily subcutaneous injections

**FOOD AND LIQUID QUALIFICATIONS:** No food restrictions (with or without food)