

RAPID ADVICE

Antiretroviral therapy for HIV infection in adults and adolescents

NOVEMBER 2009

WHO Library Cataloguing-in-Publication Data

Rapid advice: antiretroviral therapy for HIV infection in adults and adolescents

1.

ISBN

© World Health Organization 2009

All rights reserved. Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +4122 791 3264; fax: +4122 791 4857; email: bookorders@who.int). Requests for permission to reproduce or translate WHO publications - whether for sale or for noncommercial distribution - should be addressed to WHO Press, at the above address (fax: +4122 791 4806; email: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Printed in Switzerland

RAPID ADVICE

Antiretroviral therapy for HIV infection in adults and adolescents

NOVEMBER 2009

Abbreviations

| 3TC | lamivudine |
|-------------|--|
| ABC | abacavir |
| ART | antiretroviral therapy |
| ARV | antiretroviral (drug) |
| ATV | atazanavir |
| AZT | zidovudine (also known as ZDV) |
| bPI | boosted protease inhibitor |
| CD4 | T-lymphocyte bearing CD4+ receptor |
| CDC | Centers for Disease Control and Prevention |
| CIPRA HT001 | Comprehensive International Program of Research on AIDS in Haiti (clinical trial) |
| DART | Development of Antiretroviral Therapy in Africa (clinical trial) |
| d4T | stavudine |
| ddl | didanosine |
| EFV | efavirenz |
| FBC | full blood count |
| FDC | fixed-dose combination |
| FTC | emtricitabine |
| GRADE | Grading of Recommendations, Assessment, Development and Evaluation |
| НВАС | Home Based AIDS Care (clinical trial) |
| HIV | human immunodeficiency virus |
| HBV | hepatitis B virus |
| IAS | International AIDS Society |
| IRIS | immune reconstitution inflammatory syndrome |
| LPV | lopinavir |
| NNRTI | non-nucleoside reverse transcriptase inhibitor |
| NRTI | nucleoside reverse transcriptase inhibitor |
| NVP | nevirapine |
| PEPFAR | US President's Plan for Emergency AIDS Relief |
| PI | protease inhibitor |
| PMTCT | prevention of mother-to-child transmission |
| PLHIV | people living with HIV |
| /r | low-dose ritonavir |
| RCT | randomized clinical trial |
| RTI | reverse transcriptase inhibitor |
| RTV | ritonavir |
| SAPIT | Starting Antiretroviral Therapy at Three Points In Tuberculosis Therapy (clinical trial) |
| ТВ | tuberculosis |
| TDF | tenofovir disoproxil fumarate |
| VL | viral load |
| UNAIDS | Joint United Nations Programme on HIV/AIDS |
| WHO | World Health Organization |

Contents

| Abbr | Abbreviations | |
|------|--|----|
| 1. | Overview | 4 |
| 1.1 | Background | 4 |
| 1.2 | Why a revision? | 4 |
| 1.3 | Guiding principles | 4 |
| 2. | The revision process | 5 |
| 2.1 | Retrieving, summarizing and presenting the evidence | 5 |
| 2.2 | Consensus, external review and updating | 5 |
| 3. | Guideline timing and publication | 6 |
| 4. | Dissemination, adaptation, implementation and evaluation | 7 |
| 5. | Declarations of interest | 8 |
| 6. | Collaboration with external partners | 9 |
| 7. | Key Recommendations | 10 |
| 8. | References | 15 |
| 9. | Annex 1 | 17 |

1. Overview

1.1 Background

The guideline Antiretroviral therapy for hiv infection in adults and adolescents, developed by World Health Organization (WHO), was first published in 2002, simplified in 2003 and was updated in 2006. The guideline continues to follow the principles of a public health approach, aiming to optimize outcomes, including the quality of life and survival, of people living with HIV (PLHIV), and to act as a reference tool for countries to adopt and adapt according to their national circumstances.

During 2009, WHO has worked to update the guideline through a series of coordinated efforts to review and synthesize emerging evidence on when to initiate antiretroviral therapy (ART), what drug regimens to use, and the management of co-infections and treatment failure. This evidence has been assembled following systematic reviews, GRADE' profile preparation and analysis, consultations with PLHIV, cost and economic impact studies, country-level feasibility assessment, and comparisons of current country guidelines.

The groups who developed this revised guideline were: an internal WHO ART Guideline Working Group (convened in April 2009), the ART Guideline Drafting Group (convened in September 2009), the external ART Peer Review Panel (convened in September 2009), and the full ART Guideline Review Committee (convened in October 2009). The members are listed in Annex 1.

1.2 Why a revision?

Since the last guideline revision in 2006, new and compelling evidence has become available, particularly concerning the earlier start of ART. There is increasing evidence of improved survival and reduced HIV-related illnesses with the earlier initiation of antiretroviral therapy. Studies support that improved access to ART has a significant impact on the prevention of HIV transmission. Stavudine (d4T) continues to play a critical role in the scaling-up of ART in low- and middle-income countries; however, its cumulative toxicity is unacceptable to PLHIV and to many health care providers. Newer, more patient friendly but currently more expensive ART regimens are available.

The aim of the guideline is to outline standards for high quality care of PLHIV, by providing evidence-based recommendations, while considering the risks and benefits, acceptability, feasibility, cost and financial implications.

http://www.gradeworkinggroup.org/index.htm

The target audience is primarily national treatment advisory groups.

The key recommendations contained herein are released as *Rapid advice* because several countries with the highest burden of HIV infection currently are in the process of changing their national guideline for HIV treatment and care, and updating estimates for 2010 Universal Access reporting. This *Rapid advice* focuses on two key areas of the full guideline; when to start ART and what ARVs to use in adults, adolescents, pregnant women, tuberculosis (TB) and hepatitis B (HBV) co-infection.

1.3 Guiding principles

The ART Guideline Review Committee discussed and agreed upon a set of principles that should be used in developing national treatment recommendations. The principal consideration was that public health interventions should secure the greatest likelihood of survival and quality of life for the greatest numbers of PLHIV.

1. Do no harm

When introducing changes preserve access for the sickest and most in need.

2. Ensure access and equity

All clinically eligible people should be able to enter treatment services (including ART) with fair and equitable distribution of treatment services.

3. Promote quality and efficiency

Ensure delivery of the highest standards of care within a public health approach so as to achieve the greatest health impact with the optimal use of available human and financial resources.

4. Ensure sustainability

Understand the long-term consequences of change with the vision of providing continued, life-long access to ART for those in need.

In this context, the individual rights of PLHIV should not be forfeited in the course of a public health approach.

2. The revision process

2.1 Retrieving, summarizing and presenting the evidence

The PICO^{••} questions to be considered were agreed upon by the internal WHO ART Guideline Working Group in May 2009. A series of activities then were undertaken to prepare for the October 2009 guideline review meeting:

- GRADE profiles were prepared for the key PICO questions:
 when to start ART;
 - what to use in first-line and second-line antiretroviral regimens;
 - iii. when to switch to a second-line regimen.
- 2. Systematic reviews of the literature were conducted on ARV drug interactions with drugs for TB, hepatitis, malaria, and opioids; ART management for HIV/HBV co-infection; ART toxicity summaries for tenofovir, zidovudine, nevirapine and stavudine; the safety of efavirenz; the teratogenicity of efavirenz; a low-dose stavudine safety profile; and reviews of CD4 and viral load laboratory technologies.
- An impact assessment was conducted to estimate the number of PLHIV in need of treatment according to various proposed CD4 initiation thresholds.
- 4. Consultations were conducted with three organizations representing PLHIV and the findings summarized.
- 5. Costing information was prepared based on studies of procurement and production of ARVs.
- 6. A feasibility assessment was conducted for the introduction of the proposed guideline in Malawi (a similar assessment is ongoing in Tanzania).
- A report was produced on issues related to adherence to ART for PLHIV, including adolescents.
- 8. A review was undertaken to study and compare current ART guidelines from 26 countries.
- ART failure criteria were reviewed using data from ART-LINC and other studies.

The GRADE evidence profiles, and the full set of supporting documentation will be included and or referenced in the guideline.^{...}

2.2 Consensus, external review and updating

Existing recommendations were reviewed and revised at the September 2009 meeting of the ART Guideline Drafting Group. Based on presentations and individual review of the available evidence described above, the group compiled risk-benefit analysis for each area of interest. A table that includes the following domains: existing recommendations; new recommendations; quality and grade of evidence for the critical outcomes (mortality, disease progression and serious adverse events); benefits; risks; values; acceptability; costs; feasibility; gaps and research needs was prepared for each set of recommendations. Disagreements were debated during plenary and group sessions and consensus sought.

The ART Peer Review Panel reviewed and revised the riskbenefit tables and draft recommendations prepared by the ART Guideline Drafting Group and responded with their comments and other input via the ART Guidelines 2009 page of WHO's EZCollab website. Reviewers debated and exchanged ideas within EZCollab and during teleconferences. Their suggestions and comments were collated and posted on the EZCollab website. The revised recommendations were presented to the ART Guideline Review Committee in October 2009.

At the October 2009 ART Guideline Review Meeting, each subject area was reviewed, discussed and consensus sought on each recommendation and the ranking of each recommendation.

The full guideline document will be reviewed again by 2012.

^{**} PICO is an acronym that describes the elements of a well-formed clinical question. The structure includes: 'P' for the patient or population; 'I' for the intervention of interest; 'C' for comparison; and 'O' for outcome

 $^{^{\}star\star\star}$ The full set of background materials can be requested by emailing hiv-aids@ who.int.

3. Guideline timing and publication

This *Rapid advice: antiretroviral therapy for HIV infection in adults and adolescents* will be published online. Given the short life-span of the Rapid Advice, it will be posted online in English. However, requests from WHO Regional Offices for translation into other languages will be supported from WHO Headquarters.

It is expected that a draft of the full guideline will be available in February 2010 for final clearance, with publication and dissemination anticipated to start in March or April 2010.

4. Dissemination, adaptation, implementation and evaluation

WHO is working closely with WHO Regional and Country Offices, UN and other implementing partners to plan for rapid dissemination, adaptation and implementation of the new recommendations. Much experience has been obtained from previous ART guidelines, and active support for guideline revision at country level is needed. Key steps in the dissemination include:

- Release of the Rapid advice
- Production and publication of the full guideline, with translation into other languages
- Rapid development of adaptation and transition tools
- Briefings and joint planning for dissemination with international and national implementing partners
- Regional conferences and workshops, to support country adaptation

Adaptation and transition tools are designed to:

- Assist countries prioritize limited resources to facilitate full
 implementation over time
- Not compromise ART access or exclude those most in need
- Not disrupt existing scale up efforts or threaten adherence
- Move progressively towards adopting all recommendations

The WHO and UNAIDS are finalizing the global and country tools required for estimations of treatment and resource needs, in time for 2010 universal access reporting.

5. Declarations of interest

Forms were collected from every member of each group. Two declarations were made and were determined not to constitute conflicts of interest by the core guideline team. The ART Peer Review Panel was advised via the WHO's EZCollab website of a member's declaration of potential conflict (P. Cahn) and the ART Peer Review Panel concluded that there was no conflict of interest. The declaration of potential conflict of a member of the ART Guideline Review Committee (C. Holmes) was discussed with the assembled ART Guideline Review Committee at the October 2009 meeting. The ART Guideline Review Committee concluded that there was no conflict of interest.

6. Collaboration with external partners

There are no external collaborators specific to this *Rapid advice*. However, several partners have been engaged in the development of the guideline. All collaborations will be detailed in the full guideline.

Funding to support this work comes from The US President's Emergency Plan for AIDS Relief (PEPFAR), The United Nations Joint Programme on HIV/AIDS Unified Budget and Workplan (UNAIDS UBW), and specific funds through staff time.

7. Key Recommendations

RECOMMENDATION 1 When to start

- Start antiretroviral treatment in all patients with HIV who have CD4 count ≤350 cells/mm³ irrespective of clinical symptoms.
 - (Strong recommendation, moderate quality of evidence)
- CD4 testing is required to identify if patients with HIV and WHO clinical stage 1 or 2 disease need to start antiretroviral treatment.

(Strong recommendation, low quality of evidence)

 Start antiretroviral treatment in all patients with HIV and WHO clinical stage 3 or 4 irrespective of CD4 count. (Strong recommendation, low quality of evidence)

Remarks: In developing these recommendations, the panel placed high value on avoiding death, disease progression and HIV transmission over and above cost and feasibility. The recommendations are supported by moderate quality of evidence for critical patient and public health outcomes from one RCT, the CIPRA-HT001 (a singlecentre trial in Haiti) and one post hoc analysis nested in a RCT, the SMART trial (a multicentre study in 33 predominantly high income countries).^{1,2} In the GRADE profile, pooled data from these two studies provide moderate evidence that starting ART at CD4 levels higher than 200 or 250 cells/mm³ reduces mortality rates in asymptomatic, ART-naive, HIV-infected people. The panel also reviewed large observational data sets from both resource limited and high resource settings which are consistent with data from the RCTs but these did not add to the overall quality of the evidence.³⁻⁶ The panel considered that starting ART earlier is feasible if introduced in a phased manner, with the speed and completeness determined by health system capacity, HIV burden, ART coverage, equity of access and funding.

Considering the uncertain prognostic value of some WHO clinical stage 2 conditions and recent modelling and observational data suggesting that more than 50% of HIV-infected patients with this clinical stage have a CD4 count of \leq 350 cells/mm³, the panel recommended HIV-infected individuals with WHO clinical stage 1 and 2 should have access to CD4 testing to decide if treatment should be initiated.

RECOMMENDATION 2 What to start

Start one of the following regimens in ART-naïve individuals eligible for treatment. AZT + 3TC + EFV AZT + 3TC + NVP TDF + 3TC or FTC + EFV TDF + 3TC or FTC + NVP (Strong recommendation, moderate quality of evidence)

Remarks: In developing these recommendations the panel placed high value on avoiding the disfiguring, unpleasant and potentially life threatening toxicity of d4T, the need to select regimens suitable for use in most patient groups, and the benefits of using fixed dose combinations.

Current evidence suggests that these regimens are comparable in terms of efficacy, with a better overall toxicity profile than d4T based regimens. The panel was reassured by the GRADE profile evidence from RCTs, non-randomised trials and observational studies from low- middle-income countries that indicate no superiority of AZT over TDF, or NVP over EFV as part of combination ART for treatmentnaïve individuals.

On the issue of progressive reduction in the use of d4T, in settings where d4T regimens are used as the principal option for starting ART, countries should develop a plan to move towards AZT- or TDF-based first-line regimens, based on an assessment of the cost and feasibility. Systems to prevent, monitor and manage d4T-related toxicities should be implemented.

It is recommended that programs select the preferred regimen(s) applicable to the majority of PLHIV. The introduction of safer but currently more expensive first-line ARTs needs to be phased-in as currently they may not be feasible or affordable in many high-burden settings with low coverage, less developed health systems, limited laboratory capacity, finite budgets and competing health priorities. In countries with high coverage and more developed health systems, transition to new treatment regimens should occur sooner.

WHO will develop tools to assist countries/programs in the transition to and implementation of these recommendations.

RECOMMENDATION 3 ART for HIV/tuberculosis co-infection

- Start ART in all HIV-infected individuals with active tuberculosis (TB) irrespective of CD4 cell count. (Strong recommendation, low quality of evidence)
- Start TB treatment first, followed by ART as soon as possible after starting TB treatment. (Strong recommendation, moderate quality of evidence)
- 3. Use efavirenz (EFV) as the preferred non-nucleoside reverse transcriptase inhibitor (NNRTI) in patients starting ART while on TB treatment. (Strong recommendation, high quality of evidence)

Remarks: In developing these recommendations, the panel placed high value on the reduction of early mortality from HIV/TB coinfection, the reduction of TB transmission when ART is initiated earlier in all individuals with TB and improved management of TB.

On the question of when to initiate ART in TB infection, one RCT (SAPIT trial) provides moderate evidence for early initiation of ART in terms of reduced all-cause mortality, improved TB outcomes and reduced incidence of immune reconstitution inflammatory syndrome (IRIS), but there are limited data on the need to initiate ART in patients with TB and CD4 >350 cells/mm³.⁷

On the question of the impact on TB transmission and incidence, ART has been reported to reduce TB rates by up to 90% at an individual level, by 60% at a population level and to reduce TB recurrence rates by 50%.⁸⁻¹⁰ Modeling suggests that initiation of ART for all those with HIV/TB co-infection, if accompanied by high levels of coverage and ART adherence, reduces the number of TB cases, TB mortality rates and TB transmission at a population level.¹¹

Further modeling suggests that starting ART less than five years after initial HIV infection could reduce the incidence of TB by 60% to 70%. (B Williams, submitted for publication)

On the question of which NNRTI to start, the recommendations in the 2006 ART guideline were maintained, specifically EFV is recommended because of less interaction with rifampicin compared to NVP.

RECOMMENDATION 4 ART for HIV/HBV co-infection

- Start ART in all HIV/HBV co-infected individuals who require treatment for their HBV infection, irrespective of CD4 cell count or WHO clinical stage. (Strong recommendation, low quality of evidence)
- Start TDF and 3TC or FTC containing antiretroviral regimens in all HIV/HBV co-infected individuals needing treatment. (Strong recommendation, moderate quality of evidence)

Remarks: In developing these recommendations, the panel placed high value on promoting HBV diagnosis and more effective treatment of HIV/HBV co-infection.

The systematic review on this topic did not find RCTs which addressed critical HIV outcomes (death, disease progression, severe adverse events) and the GRADE profile reported only on outcomes related to HBV (HBV viral load and HBV drug resistance).

On the question of when to start ART in HIV/HBV co-infection, there are no trials comparing early versus late initiation of ART. However, observational data support that individuals with HIV/HBV co-infection have a 3- to 6-fold risk of developing chronic HBV, an increased risk of fibrosis and cirrhosis and a 17-fold increased risk of death compared to HBV-infected individuals without HIV infection. Similarly, observational data support a reduction in liver related disease with earlier and HBV-active combination ART.

On the question of what ART to start in HIV/HBV co-infection, there are data from one RCT supporting the use of at least two agents with activity against HBV in terms of improved HBV viral load response and reduced development of HBV drug resistance.¹²

RECOMMENDATION 5 ART for pregnant women

- Start ART in all pregnant women with HIV and CD4 count ≤350 cells/mm³, irrespective of clinical symptoms. (Strong recommendation, moderate quality of evidence)
- CD4 testing is required to identify if pregnant women with HIV and WHO clinical stage 1 or 2 disease need to start antiretroviral treatment or prophylaxis. (Strong recommendation, low quality of evidence)
- Start ART in all pregnant women with HIV and WHO clinical stage 3 or 4, irrespective of CD4 count. (Strong recommendation, low quality of evidence)
- 4. Start one the following regimens in ART-naïve pregnant women eligible for treatment.
 (Strong recommendation, moderate quality of evidence)
 AZT + 3TC + EFV
 AZT + 3TC + NVP
 TDF + 3TC or FTC+ EFV
 TDF + 3TC or FTC + NVP
- 5. Do not start EFV during the first-trimester of pregnancy. (Strong recommendation, low quality of evidence)

Remarks: In developing these recommendations, the ART and PMTCT panels placed high value on ensuring treatment is started early for pregnant women with HIV to avoid mother-to-child transmission and improve maternal and child-health outcomes, over and above concerns for the cost or feasibility.

On the question of **when to start**, no studies specific to pregnant women were identified. Overall evidence supports strong recommendations for the reduction of mortality, disease progression, severe adverse events, the risk of TB and the risk of HIV transmission (sexual and vertical). As with the recommendation on when to start in the general population, the panel recognised the uncertainty around the prognostic value of some WHO clinical stage 2 conditions and data from modelling and observational studies indicate that more than 50% of HIV-infected patients with this clinical stage have a CD4 count of \leq 350 cells/mm³. Therefore, the panel recommended all pregnant women with WHO clinical stages 1 and 2 have access to CD4 testing to decide when to start treatment.

On the question of **what to start**, no GRADE profiles were prepared as no RCTs were identified for the use of AZT+3TC+NVP specifically in pregnant women. Cohort studies report reduction of HIV transmission and death.¹³ There is no evidence to suggest an increase in maternal serious adverse events and no studies specifically evaluating maternal response to ART. Pregnancy registry data on the use of TDF in pregnancy show no signals to raise concern, and there is no evidence to suggest that TDF+3TC or FTC is not an acceptable alternative to AZT+3TC.¹⁴

There is very low quality, conflicting evidence on the risks of EFV causing neural tube defects. The overall rates of birth defects reported in association with EFV, NVP, lopinavir/ritonavir (LPV/r) or TDF appear similar and are consistent with rates reported in congenital defects' registries from general populations. Since neural tube closure occurs in the first 28 days and very few pregnancies are recognised by this time, the actual risks from starting EFV based ART in the first trimester are difficult to estimate.

The review of NVP safety in pregnant women with CD4 count 250–350 cells/mm³ did not confirm an increased risk of serious adverse events and the panel concluded that the benefits of using NVP in this situation outweigh the risks of not initiating ART.

The panel was unable to conclude from the evidence reviewed whether there are benefits associated with the use of EFV vis a vis NVP in pregnant women after the first trimester and with higher or unknown CD4 cell counts, although more than half of the panel members preferred EFV in these situations.

RECOMMENDATION 6 When to switch ART

- Where available, use viral load (VL) to confirm treatment failure. (Strong recommendation, low quality of evidence)
- Where routinely available, use VL every 6 months to detect viral replication. (Conditional recommendation, low quality of evidence)
- A persistent VL above 5 000 copies/ml confirms treatment failure. (Conditional recommendation, low quality of evidence)
- When VL is not available, use immunological criteria to confirm clinical failure. (Strong recommendation, moderate quality of evidence)

Remarks: In developing these recommendations, the panel was concerned by the limitations of clinical and immunological monitoring for diagnosing treatment failure, and placed high value on avoiding premature or unnecessary switching to expensive second line ART. They also valued the need to optimize the use of virological monitoring and ensure adherence.

Based on the pooled analysis of the size effects from two randomized trials (HBAC and DART trials), clinical monitoring alone (compared to combined immunological and clinical monitoring) resulted in increases in mortality, disease progression and unnecessary switches, but no difference in serious adverse events.^{15, 16} However, in one of these trials (HBAC trial), combined immunological, and clinical monitoring was compared to combined virological, immunological, and resulted in no difference in mortality, disease progression and unnecessary switches, but no difference in serious adverse events.^{15, 16} However, in one of these trials (HBAC trial), combined immunological and clinical monitoring was compared to combined virological, immunological, and clinical monitoring, and resulted in no difference in mortality, disease progression, unnecessary switches or virological treatment failures.¹⁵

RECOMMENDATION 7 Second-line ART

- A boosted protease inhibitor (PI/r) plus two nucleoside analogues (NRTIs) are recommended for second-line ART. (Strong recommendation, moderate quality of evidence)
- ATV/r and LPV/r are the preferred boosted PI's for secondline ART. (Strong recommendation, moderate quality of evidence)
 - (strong recommendation, moderate quality of evidence)
- 3. Simplification of second NRTI options is recommended.
 - If d4T or AZT has been used in first-line, use TDF + 3TC or FTC as the NRTI backbone in second-line.
 - If TDF has been used in first-line, use AZT + 3TC as the NRTI backbone in second-line.

(Strong recommendation, moderate quality of evidence)

Remarks: In developing these recommendations, the panel placed high value on using simpler second-line regimens and the availability of heat-stable, fixed-dose combinations.

On the question of whether PI monotherapy could be used as secondline ART, there is moderate quality of evidence from nine RCTs and individual study reports showing less virological suppression and higher rates of viral rebound for PI monotherapy compared to standard triple ART regimens.¹⁷⁻²² The panel concluded that an NRTI backbone should be maintained.

On the question of which bPI to use in second-line therapy, there is moderate quality of evidence that ATV/r is non-inferior to LPV/r (in combination with TDF and an optimized second NRTI) in treatment-experienced patients.²³ Non-serious adverse events varied by boosted PI with no significant difference in serious adverse events.

On the question of which NRTI backbone to use in second-line, few studies of relevance were identified. The recommended NRTI sequencing is based on likely resistance mutations and potential for retained antiviral activity. If AZT+3TC are used in first-line with sensitive monitoring and early switching, the NRTIs with remaining activity are TDF and ddl, (both very likely) and ABC (likely). In the scenario of insensitive monitoring and late switching, TDF and ddl activity are less likely, with activity of ABC unlikely. If TDF+3TC are use in first-line, with early switching, the NRTIs with remaining activity are AZT and d4T (very likely), ddl, ABC (possible). In the scenario of late switching, activity of AZT and d4T is very likely, with activity of ddl and ABC unlikely. Retained activity of 3TC is likely in the early switching scenario and less likely in the case of late switching. AZT+3TC, TDF+FTC and TDF+3TC are available as fixed-dose combinations.

RECOMMENDATION 8 Third-line regimens

- National programmes should develop policies for third-line therapy that consider funding, sustainability and the provision of equitable access to ART. (Conditional recommendation, low quality of evidence)
- Third-line regimens should include new drugs likely to have anti HIV activity such as integrase inhibitors and second generation NNRTIs and PIs. (Conditional recommendation, low quality of evidence)
- Patients on a failing second-line regimen with no new ARV options, should continue with a tolerated regimen. (Conditional recommendation, very low quality of evidence)

Remarks: The panel were concerned by unpublished cohort reports of high mortality among patients failing second-line therapy, but placed high value on balancing the need to develop policies for third-line therapy whilst maintaining increased access to first-line therapy. It was recognized that many counties have financial constraints that might limit the adoption of third-line regimens.

From a targeted literature review of relevant studies, the evidence is limited with few studies of newer agents in resource-limited settings. Data from RCTs predominantly in developed countries are available for boosted darunavir, etravirine and raltegravir. Taken together, these data support the efficacy of these agents in highly ART experienced patients. There was no uncertainty among the panel concerning the need for third-line regimens. However, there was uncertainty about how making third-line regimens available would impact the provision of first-line ART. There was also uncertainty about what third-line drugs should be provided and many studies are ongoing.

8. References

- 1. Severe P, Pape J, Fitzgerald D et al. A Randomized Clinical Trial of Early Versus Standard Antiretroviral Therapy for HIV-Infected Patients with a CD4 T Cell Count of 200-350 Cells/ml (CIPRAHT001). 49th Interscience Conference on Antimicrobial Agents and Chemotherapy; San Francisco, 2009; Abstract H-1230c.
- 2. Emery S, Neuhaus JA, Phillips AN, Babiker A, Cohen CJ, Gatell JM, et al. Major clinical outcomes in antiretroviral therapy (ART)-naive participants and in those not receiving ART at baseline in the SMART study. J Infect Dis. 2008 Apr 15;197(8):1133-44.
- 3. Moh R, Danel C, Messou E, Ouassa T, Gabillard D, Anzian A, et al. Incidence and determinants of mortality and morbidity following early antiretroviral therapy initiation in HIV-infected adults in West Africa. AIDS. 2007 Nov 30;21(18):2483-91.
- 4. Badri M, Bekker LG, Orrell C, Pitt J, Cilliers F, Wood R. Initiating highly active antiretroviral therapy in sub-Saharan Africa: an assessment of the revised World Health Organization scaling-up guidelines. AIDS. 2004 May 21;18(8):1159-68.
- 5. Erhabor O, Ejele OA, Nwauche CA. The effects of highly active antiretroviral therapy (HAART) of stavudine, lamivudine and nevirapine on the CD4 lymphocyte count of HIV-infected Africans: the Nigerian experience. Niger J Clin Pract. 2006 Dec;9(2):128-33.
- 6. Wong KH, Chan KC, Cheng KL, Chan WK, Kam KM, Lee SS. Establishing CD4 thresholds for highly active antiretroviral therapy initiation in a cohort of HIV-infected adult Chinese in Hong Kong. AIDS Patient Care STDs. 2007 Feb;21(2):106-15.
- 7. Abdool-Karim SS, Naidoo K, Grobler A, Padayatchi N, Nair G, Bamber S et al. Initiating ART during TB Treatment Significnatly Increases Survival: Results of a Randomized Controlled Clinical trial in TB/HIV co-infected Patients in South Africa. 16th Conference on Retroviruses and Opportunistic Infections; Montreal, 2009. Oral abstract 36a. [Available from http://www. retroconference.org/Abstracts/34255.htm]
- 8. Lawn SD, Churchyard G. Epidemiology of HIV-associated tuberculosis. Curr Opin HIV AIDS. 2009 Jul;4(4):325-33.
- 9. Middelkoop K, Wood R, Myer L, Sebastian E, Bekker LG. Can antiretoviral therapy contain a previously escalating TB epidemic in a high HIV prevalence community. 5th IAS Conference on HIV Pathogenesis, Treatment and Prevention; 2009; Cape Town. Abstract CDB 041 [Available from http://www.ias2009.org/pag/Abstracts.aspx?AID=2932]
- Golub JE, Astemborski J, Ahmed M, Cronin W, Mehta SH, Kirk GD, et al. Long-term effectiveness of diagnosing and treating latent tuberculosis infection in a cohort of HIV-infected and at-risk injection drug users. J Acquir Immune Defic Syndr. 2008 Dec 15;49(5):532-7.
- 11. Atun RA, Lebcir RM, Drobniewski F, McKee M, Coker RJ. High coverage with HAART is required to substantially reduce the number of deaths from tuberculosis: system dynamics simulation. Int J STD AIDS. 2007 Apr;18(4):267-73.
- 12. Matthews GV, Avihingsanon A, Lewin SR, Amin J, Rerknimitr R, Petcharapirat P, et al. A randomized trial of combination hepatitis B therapy in HIV/HBV coinfected antiretroviral naive individuals in Thailand. Hepatology. 2008 Oct;48(4):1062-9.
- 13. Bae WH, Wester C, Smeaton LM, Shapiro RL, Lockman S, Onyait K, et al. Hematologic and hepatic toxicities associated with antenatal and postnatal exposure to maternal highly active antiretroviral therapy among infants. AIDS. 2008 Aug 20;22(13):1633-40.
- 14. The antiretroviral pregnancy registry [database on the Internet]. Antiretroviral Pregnancy Registry Steering Committee. 2009. [Available from http://www.apregistry.com/forms/interim_report.pdf]
- 15. Coutinho A, Mermin J, Ekwaru J, Were W, R Bunnell, Kaharuza F et al. Utility of Routine Viral Load, CD4 Cell Count, and Clinical Monitoring among HIV-infected Adults in Uganda: A Randomized Trial. 15th Conference on Retroviruses and Opportunistic Infections; 2008; Boston, Abstract 125. [Available from http://www.retroconference.org/2008/ Abstracts/30881.htm]

- 16. Mugyenyi P, Walker S Hakim J, Munderi P, Gibb D, Kityo C, et al. Impact of routine laboratory monitoring over 5 years after antiretroviral therapy (ART) initiation on clinical disease progression of HIV-infected African adults: the DART trial final results 2009. 5th IAS Conference on HIV Pathogenesis, Treatment and Prevention; 2009; Cape Town. Abstract TUSS 102 [Available from http://www.ias2009.org/pag/Abstracts.aspx?AID=3807]
- 17. Arribas JR, Pulido F, Delgado R, Lorenzo A, Miralles P, Arranz A, et al. Lopinavir/ritonavir as single-drug therapy for maintenance of HIV-1 viral suppression: 48-week results of a randomized, controlled, open-label, proof-of-concept pilot clinical trial (OK Study). J Acquir Immune Defic Syndr. 2005 Nov 1;40(3):280-7.
- 18. Escobar I, Pulido F, Perez E, Arribas JR, Garcia MP, Hernando A. [Pharmacoeconomic analysis of a maintenance strategy with lopinavir/ritonavir monotherapy in HIV-infected patients]. Enferm Infecc Microbiol Clin. 2006 Oct;24(8):490-4.
- Cameron DW, da Silva Barbara A, Arribas Jose R, Myers Robert A, Bellos Nicholaos C, Gilmore N, et al. A 96 Week Comparison of Lopinavir Ritonavir Combination Therapy Followed by Lopinavir Ritonavir Monotherapy versus Efavirenz Combination Therapy. J Infect Dis. 2008;198(2):234-40.
- 20. J. Arribas AH, J. Gerstoft, G. Fatkenheuer, M. Nelson, N. Clumeck, F. Pulido et al. The MONET trial: darunavir/ritonavir monotherapy shows non-inferior efficacy to standard HAART, for patients with HIV RNA < 50 copies/mL at baseline. 5th IAS Conference on HIV Pathogenesis, Treatment and Prevention; 2009; Cape Town. Abstract .TUAB 106-LB [Available from http:// www.ias2009.org/pag/Abstracts.aspx?AID=3626]
- Katlama C, Valentin MA, Algarte-Genin M, Duvivier C, Lambert-Niclot S, Girard PM et al. Efficacy of darunavir/ritonavir as single-drug maintenance therapy in patients with HIV-1 viral suppression: a randomized open-label non-inferiority trial, MONOI-ANRS 136. 5th IAS Conference on HIV Pathogenesis, Treatment and Prevention; 2009; Cape Town. Abstract WELBB 102. [Available from http://www.ias2009.org/pag/Abstracts.aspx?AID=3689]
- 22. Delfraissy JF, Flandre P, Delaugerre C, Ghosn J, Horban A, Girard PM, et al. Lopinavir/ritonavir monotherapy or plus zidovudine and lamivudine in antiretroviral-naive HIV-infected patients. AIDS. 2008 Jan 30;22(3):385-93.
- 23. Molina JM, Andrade-Villanueva J, Echevarria J, Chetchotisakd P, Corral J, David N, et al. Once-daily atazanavir/ritonavir versus twice-daily lopinavir/ritonavir, each in combination with tenofovir and emtricitabine, for management of antiretroviral-naive HIV-1-infected patients: 48 week efficacy and safety results of the CASTLE study. Lancet. 2008 Aug 23;372(9639):646-55.

9. Annex 1

WORLD HEALTH ORGANIZATION

2009 ART Guideline Review Meeting

Hotel Ramada Encore, Geneva, 14-16 October 2009

LIST OF PARTICIPANTS

ART Guideline Review Committee

Lori Bollinger (Health Economist) Vice President Futures Institute 41-A New London Turnpike Glastonbury, CT 06033 USA Ibollinger@futuresinstitute.org

Alexandra Calmy (Hiv Researcher/Clinician)

Cheffe de clinique scientifique Service de Maladies Infectieuses Unité VIH/SIDA Université de Genève SWITZERLAND alexandra.calmy@hcuge.ch

Larry William Chang (Grade Methodologist)

Johns Hopkins School of Medicine 1503 E. Jefferson St. Baltimore, MD 21287 USA larrywillchang@gmail.com

Zengani Chirwa (Implementing Partner)

ART Technical Advisor Ministry of Health, Lilongwe MALAWI czenga@hotmail.com

Francois Dabis

(HIV Researcher/Epidemiologist) Centre de Recherche INSERM U.897 Institut de Santé Publique, Epidémiologie et Développement (ISPED) Université Victor Segalen Bordeaux 2 France francois.dabis@isped.u-bordeaux2.fr

Mathias Egger (HIV Researcher/Epidemiologist)

Professor of Epidemiology & Public Health Institute of Social & Preventive Medicine University of Bern Finkenhubelweg 11 CH-3012 Bern, SWITZELAND egger@ispm.unibe.ch

Shaffiq Essajee (Implementing Partner)

Director of Clinical Operations Senior Advisor in Pediatrics Clinton Foundation HIV/AIDS Initiative USA sessajee@clintonfoundation.org

Loon Gangte (PLHIV representative)

President, Delhi, Network of Positive People (DNP+) Gali No.3,House No-64 Neb Sarai, New Delhi-1100 68 INDIA Ioon_gangte@yahoo.com

Julian Gold (Hiv Researcher/Clinician)

Director, Albion St Centre 150 Albion St. Surry Hills NSW 2010 Australia julian.gold@sesiahs.health.nsw.gov.au

Sarah Glover (HIV Researcher)

London School of Hygiene and Tropical Medecine Keppel St London, WC1E 7HT UNITED KINGDOM glover_sarah@yahoo.com

James Hakim (Chair/Hiv Researcher/Clinician) Professor University of Zimbabwe Harare ZIMBABWE jhakim@mweb.co.zw

Charles Holmes (Implementing Partner)

Senior Technical Advisor Office of the US Global AIDS Coordinator S/GAC, SA-29, 2nd Floor Washington, DC 20522-2920 USA holmescb@state.gov

Elizabeth Humphreys (GRADE methodologist)

Cochrane Collaborative Group on HIV/AIDS University of California San Francisco USA elizahumphreys@gmail.com

Robert Mwanri Josiah (Implementing Partner)

Head, Care And Treatment Unit National Aids Control Programme (NACP) Luthuli Road, P.O. BOX 11875 Dar Es Salaam TANZANIA rmwjosiah@gmail.com

Ayesha Khan (Programme Implementer)

Infectious Diseases and M&E Specialist Health Systems Strengthening and Policy Unit (HSSPU) Ministry of Health Health Services Academy Chak Shahzad, Islamabad PAKISTAN ayakhan@gmail.com

Stephen Lawn (TB/HIV Researcher)

Reader in Infectious Diseases & Tropical Medicine Desmond Tutu HIV Centre, IIDMM, University of Cape Town, Anzio Road, Observatory, 9758 Cape Town SOUTH AFRICA stevelawn@yahoo.co.uk

Jean-Paul Moatti (Health Economist) UMR INSERM/IRD 912

ORS PACA 23 rue Stanislas Torrents 13006 MARSEILLE FRANCE jean-paul.moatti@inserm.fr

Lynne Mofenson (HIV Researcher/Clinician)

Pediatric, Adolescent and Maternal AIDS Branch Center for Research for Mothers and Children Eunice Kennedy Shriver National Institute of Child Health and Human Development National Institutes of Health 6100 Executive Boulevard, Room 4B11 Rockville, MD 20852 USA Im65d@nih.gov

Paula Munderi (HIV Researcher/Clinician)

Medical Research Council Programme on AIDS Uganda Virus Research Institute PO Box 49 Entebbe UGANDA paula.munderi@mrcuganda.org

Irene Mukui (Programme Implementer)

Ministry of Public Health and Sanitation Department of Diseases Prevention and Control Kenyanta Hospital Ground P. O. Box 19982-00202 Nairobi KENYA njahira2004@yahoo.com

Mutinta Nalubamba-Phiri (Programme Implementer)

Paediatric ART Programme Officer P.O.Box 32588, Lusaka ZAMBIA drtinta@yahoo.com

Portia Ngcaba (PLHIV Representative)

TAC Eastern Cape Provincial Coordinator NGO Sector Treatment Action Campaign 57 a Paterson Street Arcadia East London SOUTH AFRICA portia@mail.tac.org.za

Sylvia Ojoo (Implementing Partner)

Senior Medical Technical Adviser, AIDSRelief Institute of Human Virology of the University of Maryland, School of Medicine KREP Centre, Wood Avenue, Kilimani PO Box 495, 00606 Nairobi KENYA sylvia_ojoo@yahoo.co.uk

Vladimir Osin (PLHIV representative)

Public Relations Manager Psychological centre "Dvizhenie" Moscow RUSSIA osin.vladimir@gmail.com

Praphan Phanuphak

(HIV researcher/clinician/implementing partner) Director The Thai Red Cross AIDS Research Centre Co-Director, HIV-NAT 104 Rajdamri Road Bangkok 10330 THAILAND praphan.p@chula.ac.th

Bharat Bhushan Rewari

(HIV clinician/implementing partner) Sr.Physician, Dr RML Hospital& National Programme Officer (ART) National AIDS Control Organisation, 6th Floor, Chandralok Building, 36, Janpath, New Delhi-110001 INDIA drbbrewari@yahoo.com

George Rutherford (Grade Methodologist)

University of California, San Francisco Institute for Global Health 50 Beale Street, Suite 1200 San Francisco, California 94105 USA grutherford@psg.ucsf.edu

Papa Salif Sow (HIV researcher/clinician)

Dakar University Hospital Fann Hospital BP 5035 Dakar SENEGAL salifsow@orange.sn

Erik Schouten (implementing partner)

Technical Assistant for Co-ordination of HIV and AIDS Programmes Department of HIV and AIDS Ministry of Health P.O. Box 30377 Lilongwe MALAWI eschouten.mw@gmail.com

Tengiz Tsertsvadze (Programme Implementer)

Director General Infectious Diseases AIDS and Clinical Immunology Research Center National AIDS Coordinator Tbilisi State University faculty of Medicine Tbilisi, GEORGIA tengizt@gol.ge

Rochelle Walensky (HIV researcher/health economist) Harvard Center for AIDS Research Massachusetts General Hospital 50 Staniford Street, 9th floor Boston, MA 02114 USA rwalensky@partners.org

Robin Wood (HIV researcher/clinician) (UCT), Faculty of Health 26 Orchand Heights, Newlands, Cape Town SOUTH AFRICA Email: robin.wood@hiv-research.org.za

Augustin Okenge Yuma (Programme Implementer) Directeur PNLS Programme National de Lutte contre le VIH/SIDA et les ISTTel Kinshasa DEMOCRATIC REPUBLIC OF CONGO dokenge@yahoo.fr

Oleg Yurin (HIV researcher/epidemiologist) Deputy Head of Federal AYDS Centre, Leading Researcher of Central Institute of Epidemiology/ Moscow RUSSIA Email: oleg_gerald@mail.ru

Megan O'Brien (HIV researcher) Clinton Foundation HIV/AIDS Initiative 55 West 125th Street New York, NY10027 USA mobrien@clintonfoundation.org

Suzanne Crowe (HIV researcher/laboratory) Head, Centre for Virology Burnet Institute 85 Commercial Road, Melbourne, Victoria, Australia 3004 GPO Box 2284, Melbourne, Victoria, AUSTRALIA crowe@burnet.edu.au

Elliot Raizes (observer) (implementing partner) Care and Treatment Branch NCHSTP/GAP CDC MS-E04 1600 Clifton Road NE Atlanta, GA 3033 USA gwq0@cdc.gov

20

WHO Staff

Shelley Agarwall Intern Antiretroviral Treatment and HIV Care

Silvia Bertagniolo Medical Officer Antiretroviral Treatment and HIV Care

Txema Calleja Medical Officer Strategic Information

Siobhan Crowley Acting Coordinator Antiretroviral Treatment and HIV Care

Micheline Diepart Medical Officer Antiretroviral Treatment and HIV Care

Boniface Dongbo Technical Officer Systems Strengthening and HIV

Chris Duncombe Consultant Antiretroviral Treatment and HIV Care

Sally Girvin Consultant Antiretroviral Treatment and HIV Care

Reuben Granich Medical Officer Anti-retroviral Treatment and HIV Care

Teguest Guerma Acting Director HIV/AIDS Department

Frank Lule AFRO Jacqueline Ying-Ru Lo Coordinator Prevention in the Health Sector

Neil Parkin Medical Officer Anti-retroviral Treatment and HIV Care

Francoise Renaud-Thery Medical Officer Systems Strengthening and HIV

Nathan Shaffer Medical Officer Prevention in the Health Sector

Tin Tin Sint Medical Officer Prevention in the Health Sector

Omar Sued HIV Treatment and Care PAHO HIV/STI Project (FCH/HI) 525 23rd St. NW Washington, DC 20037 USA

Yves Souteyrand Coordinator Strategic Information

Marco Vitoria Medical Officer Antiretroviral Treatment and HIV Care

Dharani Yerrakalva Intern Antiretroviral Treatment and HIV Care

Stop TB

Haileyesus Getahun Medical Officer TB/HIV and Drug Resistance

Delphine Sculier Medical Officer TB/HIV and Drug Resistance

UNAIDS

Carlos Avila Adviser Al/RTN Resource Tracking, Needs and Costing Unit

Anja Grujovic Technical Officer AI/AFR AIDS Financing & Economic Division

ART Peer Review Panel 2009 revision of Antiretroviral therapy for HIV infection in adults and adolescents

Xavier Anglaret (HIV researcher/epidemiologist) Senior Researcher INSERM U897 Université Bordeaux 2 33076 Bordeaux Cedex FRANCE

Xavier.Anglaret@isped.u-bordeaux2.fr

Veronique Bortolotti

(programme implementer) Medical officer, HIV care and treatment and PMTCT WHO EMRO AIDS and Sexually Transmitted Diseases (ASD) Unit World Health Organization Regional Office of the Eastern Mediterranean Abdul Razzak AI-Sanhouri St. P.O. Box 7608 Nasr City Cairo 11371 EGYPT Bortolottiv@emro.who.int

Pedro Cahn (HIV researcher/clinician)

Director Fundacion Huesped Angel Peluffo 3932 C1202ABB - Buenos Aires ARGENTINA pcahn@huesped.org.ar

Serge Paul Eholié (HIV researcher/clinician)

Unit of Infectious Diseases Treichville University Hospital BPV3 CHU Treichville Abidjan, IVORY COAST speholie@afnet.net

Jean-Baptiste Guiard-Schmid (programme implementer)

Inter Country Adviser World Health Organization Inter-country Support Team - West Africa 03 BP 7019 Ouagadougou03 BuURKINA FASO guiardschmidjb@bf.afro.who.int

Mark Harrington (HIV/AIDS policy)

Executive Director Treatment Action Group 611 Broadway, Suite 308 New York, NY 10012 USA markhar@gmail.com

Elly Katabira (HIV researcher/clinician)

Professor Makerere Medical School Department of Medicine School of Medicine Makerere University College of Health Sciences P. O. Box 7072 Kampala UGANDA katabira@imul.com

Ricardo Kuchenbecker (HIV researcher/clinician)

Professor of Medicine Federal University of Rio Grande do Sul Hospital de Clínicas de Porto Alegre Rua Ramiro Barcelos, 2350 90035-903Porto Alegre, RS BRAZIL rkuchen@gmail.com

Barbara Marston (implementing partner) Medical Officer

Centers for Disease Control and Prevention 1600 Clifton Road Atlanta GA 30333 MS-E-04 USA bmarston@cdc.gov

Padmini Srikantiah (Programme implementer) Medical Officer

WHO-SEARO World Health House Indraprastha Estate Mahatma Gandhi Marg New Delhi 110001 INDIA srikantiahp@searo.who.int

Scott Hammer (HIV researcher/clinician)

Chief, Division of Infectious Diseases Columbia University 630 West 168th Street PH 8 West, Room 876 New York, New York 10032 USA smh48@columbia.edu

ART Guideline Drafting Group 2009 revision of Antiretroviral therapy for HIV infection in adults and adolescents

Dr Alexandra Calmy (HIV researcher/clinician) Cheffe de clinique scientifique Service de Maladies Infectieuses Unité VIH/SIDA Université de Genève SWITZERLAND

Larry William Chang (Grade Methodologist)

Johns Hopkins School of Medicine 1503 E. Jefferson St. Baltimore, MD 21287 USA larrywillchang@gmail.com

Dr Tomas Finkbeiner

(HIV clinician/implementing partner) Director of Clinical Services CDC Dar Es Salaam TANZANIA

Professor James Hakim

(Chair, Guideline Panel, HIV researcher/clinician) Professor University of Zimbabwe Harare ZIMBABWE

Dr Elizabeth H. Humphreys (GRADE methodologist) Cochrane Collaborative Group on HIV/AIDS University of California, San Francisco USA Dr Elliot Raizes (Implementing partner) Care and Treatment Branch NCHSTP/GAP CDC MS-E04 1600 Clifton Road NE Atlanta, GA 3033 USA

Dr. Paul J. Weidle (HIV researcher/epidemiologist) CAPT, USPHS Research Support Officer Epidemiology Branch Division of HIV/AIDS Prevention National Center for HIV, Viral Hepatitis, STD, and TB Prevention Centers for Disease Control and Prevention 1600 Clifton Rd, MS E-45 Atlanta, GA 30333 USA Dr Reuben Granich Medical Officer Anti-retroviral Treatment and HIV Care

Dr Kerry Kutch External Relations Officer HIV/AIDS Department

Ms Sally Girvin Consultant Antiretroviral Treatment and HIV Care

Dr Tin Tin Sint Medical Officer Prevention in the Health Sector

Dr Marco Vitoria Medical Officer Antiretroviral Treatment

UNAIDS Staff

Dr Carlos AVILA-FIGUEROA AI/RTN Resource Tracking, Needs & Costing Unit UNAIDS

WHO Staff

Dr Siobhan Crowley Acting Coordinator Anti-retroviral Treatment and HIV Care

Dr Chris Duncombe Consultant Antiretroviral Treatment and HIV Care

Dr Jesus Maria Garcia Calleja Medical Officer Strategic Information

Mrs Françoise Renaud-Thery Medical Officer Systems Strengthening and HIV