AIDS Trends



National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention

Division of HIV/AIDS Prevention

AIDS Diagnoses, Deaths, and Persons Living with AIDS, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Death may be due to any cause.



 \bigcirc

AIDS Diagnoses and Deaths of Adults and Adolescents with AIDS, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Death may be due to any cause.



AIDS Diagnoses among Adults and Adolescents, by Race/Ethnicity, 1985–2010—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. ^a Includes Asian/Pacific Islander legacy cases. ^b Hispanics/Latinos can be of any race.



AIDS Diagnoses among Adults and Adolescents, by Race/Ethnicity 1985–2010—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. ^a Includes Asian/Pacific Islander legacy cases.



 \supset

Deaths of Adults and Adolescents with AIDS, by Race/Ethnicity, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Death may be due to any cause.
^a Includes Asian/Pacific Islander legacy cases.
^b Hispanics/Latinos can be of any race.



Deaths of Adults and Adolescents with AIDS, by Race/Ethnicity, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Death may be due to any cause. a Includes Asian/Pacific Islander legacy cases.



AIDS Diagnoses and Deaths of Persons with AIDS, with HIV infection Attributed to Male-to-Male Sexual Contact, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Death may be due to any cause.



AIDS Diagnoses among Persons with HIV Infection Attributed to Male-to-Male Sexual Contact, by Race/Ethnicity, 1985–2010— United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. ^a Includes Asian/Pacific Islander legacy cases. ^b Hispanics/Latinos can be of any race.



AIDS Diagnoses among Persons with HIV Infection Attributed to Male-to-Male Sexual Contact, by Race/Ethnicity, 1985–2010— United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. ^a Includes Asian/Pacific Islander legacy cases.



 \supset

AIDS Diagnoses and Deaths of Persons with AIDS, with HIV Infection Attributed to Injection Drug Use, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. Death may be due to any cause.



AIDS Diagnoses and Deaths of Persons with AIDS, with HIV Infection Attributed to Heterosexual Contact^a, 1985–2009— United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. Death may be due to any cause. ^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



AIDS Diagnoses and Deaths of Persons with AIDS, with HIV Infection Attributed to Male-to-Male Sexual Contact and Injection Drug Use, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. Death may be due to any cause.



AIDS Diagnoses among Adult and Adolescent Females, with HIV Infection Attributed to Heterosexual Contact^a, by Race/Ethnicity, 1985–2010—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting.
^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.
^b Includes Asian/Pacific Islander legacy cases.
^c Hispanics/Latinos can be of any race.



Deaths of Adult and Adolescent Females with AIDS, with HIV Infection Attributed to Heterosexual Contact^a, by Race/Ethnicity, 1985–2009, United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. Death may be due to any cause.
^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.
^b Includes Asian/Pacific Islander legacy cases.
^c Hispanics/Latinos can be of any race.



Adults and Adolescents Living with an AIDS Diagnosis, by Transmission Category, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. ^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection. ^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Adult and Adolescent Males Living with an AIDS Diagnosis, by Transmission Category, 1985–2009—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. ^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection. ^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Adult and Adolescent Females Living with an AIDS Diagnosis, by Transmission Category, 1985–2009— United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays and missing risk-factor information, but not for incomplete reporting. ^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection. ^b Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



 \supset

AIDS Diagnoses among Perinatally Infected Persons, 1985–2010—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



 \bigcirc