

Table 2-2: Prevalence of Current Symptoms in Different Patient Populations with HIV Infection

PATIENT POPULATIONS					
	Moss (U.K.) (n=100) % (rank)	Foley (Canada) (n=100) % (rank)	Fantoni (Italy) (n=1128) % (rank)	LaRue (France) (n=314) % (rank)	Kelleher (U.K.) (n=118) % (rank)
Weight loss	85%(1)∇	91%(1)§	—	31%(6)	58%(1)
Anorexia	44%(3)	—	34%(2)	—	38%(3)
Pain	71%(2)	63%(3)	29%(4)	52%(1)	55%(2)
Cough	30%(5)	34%(8)	32%(3)	27%(8)	19%(6)
Dyspnea / resp. symptoms	17%(11)	48%(5)	19%(6)	22%(12)	15%(9)
Fatigue	—**	77%(2)	55%(1)	50%(2)	—
Nausea / vomiting	24%(8)	35%(7)	22%(5)	28%(7)	17%(8)
Cognitive dysfunction	32%(4)	43%(6)	—	—	19%(7)
Depression	26%(7)	32%(9)*	—	26%(10)	10%(12)*
Anxiety	—	—	—	40%(3)	—
Diarrhea	23%(9)	—	11%(8)	24%(11)	14%(10)
Mouth sores	—	—	—	33%(5)	—
Fever / sweats	—	—	—	27%(9)	—
Pruritis / dry skin	27%(6)	24%(11)	17%(7)	23%(12)	—
Constipation	22%(10)	23%(12)	—	—	25%(4)
Visual loss	—	25%(10)	12%(9)	—	—
Headache	11%(12)	—	—	—	11%(11)
Hemiparesis / ataxia	—	—	—	—	23%(5)
Incontinence	—	55%(4)	—	—	—
Insomnia	—	—	—	37%(4)	—

∇ Includes fatigue

§ Includes anorexia

* Includes anxiety

** Indicates that the symptom was not included as a separate category for this study population